Multiplicative induction and units for the ring of monomial representations

著者	Takegahara Yugen
journal or publication title	Advances in Mathematics
volume	355
year	2019-10-15
URL	http://hdl.handle.net/10258/00010040

doi: info:doi/10.1016/j.aim.2019.106768

Multiplicative induction and units for the ring of monomial representations

Yugen Takegahara*

Muroran Institute of Technology, 27-1 Mizumoto, Muroran 050-8585, Japan
E-mail: yugen@mmm.muroran-it.ac.jp

Abstract

Let G be a finite group, and let A be a finite abelian G -group. For each subgroup H of G , $\Omega(H, A)$ denotes the ring of monomial representations of H with coefficients in A , which is a generalization of the Burnside ring $\Omega(H)$ of H . We research the multiplicative induction map $\Omega(H, A) \rightarrow \Omega(G, A)$ derived from the tensor induction map $\Omega(H) \rightarrow \Omega(G)$, and also research the unit group of $\Omega(G, A)$. The results are explained in terms of the first cohomology groups $H^1(K, A)$ for $K \leq G$. We see that tensor induction for 1-cocycles plays a crucial role in a description of multiplicative induction. The unit group of $\Omega(G, A)$ is identified as a finitely generated abelian group. We especially study the group of torsion units of $\Omega(G, A)$, and study the unit group of $\Omega(G)$ as well.

Contents

1	Introduction	2
2	Monomial Burnside rings	4
2A	1-cocycles	4
2B	Isomorphism classes	6
2C	Tensor product	9
3	Multiplicative induction	12
3A	Tensor induction	12
3B	Tensor induction for 1-cocycles	16
3C	Algebraic maps	18
4	The mark homomorphism	21
4A	The first cohomology group	21
4B	The ghost ring	23
4C	Invariant of tensor induction	26

*This work was supported by JSPS KAKENHI Grant Number JP16K05052.

2010 *Mathematics Subject Classification*. Primary 19A22; Secondary 20J06, 16U60.

Keywords. Burnside ring, cocycle, idempotent, monomial Burnside ring, multiplicative induction, tensor induction, unit.

5	Fundamentals of monomial Burnside rings	30
5A	The Burnside homomorphism	30
5B	The Cauchy-Frobenius homomorphism	32
5C	Idempotents of Burnside rings	33
6	Units of Burnside rings	37
6A	The Yoshida criterion for the units of Burnside rings	37
6B	Structure of the unit groups of Burnside rings	40
7	Units of monomial Burnside rings	43
7A	The unit groups of monomial Burnside rings	43
7B	Torsion units of monomial Burnside rings	44

1 Introduction

Let G be a finite group, and let A be a finite abelian group on which G acts via a homomorphism from G to the group of automorphisms of A . We are concerned with the ring $\Omega(G, A)$ of monomial representations of G with coefficients in A , which was introduced by Dress [12] and is called the monomial Burnside ring for short. This ring contains the ordinary Burnside ring $\Omega(G)$ as a subring, and is applicable to the representation theory of finite groups. There are some well-known facts about $\Omega(G, A)$ (see, *e.g.*, [2, 3, 12, 13, 22, 23]). Many properties of Burnside rings seem to be extended to monomial Burnside rings; for instance, the prime ideal spectrum of $\Omega(G, A)$ was studied in [12] (see also [10]). In this paper, among others, we focus our mind on the concept of multiplicative induction for monomial Burnside rings and the unit group of $\Omega(G, A)$. There are some specific characterizations of them which mean the algebraic peculiarities of $\Omega(G, A)$.

Following [12], we give the concept of (G, A) -sets and define simple (G, A) -sets $(G/K)_\nu$ for $K \leq G$ and 1-cocycles $\nu : K \rightarrow A$ in Section 2. The monomial Burnside ring $\Omega(G, A)$, which is defined to be the Grothendieck ring of the category of (G, A) -sets (see Definition 2.13), is the commutative unital ring consisting of all formal $\mathbb{Z}$ -linear combinations of the symbols $[(G/K)_\nu]$ corresponding to the isomorphism classes of (G, A) -sets containing simple (G, A) -sets $(G/K)_\nu$ (see Proposition 2.14).

The concept of multiplicative induction for Burnside rings was introduced by tom Dieck [9] and Dress [11], and was developed by Yoshida [32]. In an attempt to introduce multiplicative induction for monomial Burnside rings, Barker [2] successfully defined the tenduction map $\mathbb{Z}_C \text{ten}_H^G : B(C, H) \rightarrow B(C, G)$ for each $H \leq G$, where C is a supercyclic group and $B(C, H)$ is the monomial Burnside ring for H with fibre group C , as a generalization of multiplicative induction for Burnside rings. (If C is a finite cyclic group on which G acts trivially, then $\Omega(G, C) \simeq B(C, G)$.)

In Section 3, we introduce the multiplicative induction map

$$\overline{\text{Map}}_H(G, -) : \Omega(H, A) \rightarrow \Omega(G, A), \quad x \mapsto \overline{\text{Map}}_H(G, x)$$

for each $H \leq G$. When A is a cyclic group on which G acts trivially, this map is associated with tensor induction for linear characters of G (cf. [8, §13A]). We have $\overline{\text{Map}}_H(G, [(H/H)_\sigma]) = [\widehat{\text{Map}}_H(G, (H/H)_\sigma)] = [(G/G)_{\sigma^{\otimes G}}]$ for all 1-cocycles $\sigma : H \rightarrow A$ (see Example 3.13), where 1-cocycles $\sigma^{\otimes G} : G \rightarrow A$ are obtained from $\sigma : H \rightarrow A$ by tensor induction. There is a nice formula of multiplicative (tensor) induction for Burnside rings (cf. [8, (80.49) Corollary]). The methods used in [8, §80C] enable us to establish that for any (H, A) -sets T_0 and T ,

$$\overline{\text{Map}}_H(G, [T_0] - [T]) = \sum_{i=0}^n (-1)^i [\widehat{\text{Map}}_H(G, T_0, T_1, \dots, T_i)], \quad (1.1)$$

where $n = |G : H|$ and $T = T_1 = \dots = T_n$ (see Proposition 3.22).

The mark homomorphism ρ_G , which was introduced by Dress [12], is a ring monomorphism from $\Omega(G, A)$ to the set $\mathfrak{U}(G, A) := (\prod_{K \leq G} \mathbb{Z}H^1(K, A))^G$ of G -invariants in the direct product of integral group rings of the first cohomology groups $H^1(K, A)$ for $K \leq G$, where the action of G on $\prod_{K \leq G} \mathbb{Z}H^1(K, A)$ is given by the conjugation maps $\text{con}_K^g : \mathbb{Z}H^1(K, A) \rightarrow \mathbb{Z}H^1({}^gK, A)$ for $K \leq G$ and $g \in G$. For each $U \leq G$, there is a ring homomorphism $-\otimes^G : \mathbb{Z}H^1(U, A) \rightarrow \mathbb{Z}H^1(G, A)$ derived from tensor induction which assigns to a 1-cocycle $\tau : U \rightarrow A$ the 1-cocycle $\tau^{\otimes G} : G \rightarrow A$. In Section 4, we describe $\overline{\text{Map}}_H(G, x) \in \Omega(G, A)$ for each $x \in \Omega(H, A)$ via ρ_G as

$$\rho_G(\overline{\text{Map}}_H(G, x)) = \left(\prod_{KgH \in K \backslash G/H} \text{con}_{K^g \cap H}^g (x_{K^g \cap H})^{\otimes K} \right)_{K \leq G} \in \mathfrak{U}(G, A) \quad (1.2)$$

under the assumption that $\rho_H(x) = (x_L)_{L \leq H}$, where $\rho_H : \Omega(H, A) \rightarrow \mathfrak{U}(H, A)$ is the mark homomorphism (see Theorem 4.16). This fact is a generalization of [32, §3(b.3)]. We make use of Eq.(1.1) to prove Eq.(1.2).

The fundamental theorem of the Burnside ring $\Omega(G)$ (cf. [32, Lemma 2.1]) is a useful instrument for finding the idempotents of $\Omega(G)$ (cf. [33, 4.12 Theorem]), and is also essential to the Yoshida criterion (see Theorem 6.4) for the units of $\Omega(G)$. In Section 5, we insist on the existence of a short exact sequence

$$0 \longrightarrow \Omega(G, A) \xrightarrow{\varphi} \widetilde{\Omega}(G, A) \xrightarrow{\psi} \text{Obs}(G, A) \longrightarrow 0$$

of additive groups (see Theorem 5.9) derived from the Cauchy-Frobenius lemma (see, *e.g.*, [33, 2.7 Lemma]), which generalizes the fundamental theorem of $\Omega(G)$.

Information of the primitive idempotents of the Burnside algebra $\mathbb{Q} \otimes_{\mathbb{Z}} \Omega(G)$ can help us to realize the units of $\Omega(G)$. Following [33, §4], we review the primitive idempotents of $\mathbb{Q} \otimes_{\mathbb{Z}} \Omega(G)$ and those of $\Omega(G)$; the latter are precisely the primitive idempotents of $\Omega(G, A)$ (see Theorem 5.18).

The unit group $\Omega(G)^\times$ of the Burnside ring $\Omega(G)$ is studied in many papers (see, *e.g.*, [6, 9, 11, 15, 18, 19, 20, 24, 30, 32]). Section 6 is devoted to a review of some

well-known facts about $\Omega(G)^\times$. We also study a certain specific type of units (see Proposition 6.11), and present an additional fact about the structure of $\Omega(G)^\times$ for which the Yoshida criterion plays a crucial role (see Corollary 6.18).

The unit group $\Omega(G, A)^\times$ of the monomial Burnside ring $\Omega(G, A)$ was studied in [2, 22]. In Section 7, we show that $\Omega(G, A)^\times$ is a finitely generated abelian group (see Proposition 7.2). Consequently, the group $\Omega(G, A)^\omega$ of torsion units of $\Omega(G, A)$ is a finite abelian group. The basic structure of $\Omega(G, A)^\omega$ is analyzed on the basis of a generalization of the Yoshida criterion (see Theorem 7.3). We adapt the methods presented in [2, §8] for an analysis of $\Omega(G, A)^\omega$, and successfully elucidate the structure of $\Omega(G, A)^\omega$ in the sequel (see Corollary 7.4). Specifically, if G is nilpotent, then the universal result deduces that

$$\Omega(G, A)^\omega \simeq \Omega(G)^\times \times H^1(G, A)$$

(see Example 7.6). This fact is a generalization of [22, Proposition 5.1].

Notation Let G be a finite group. We denote by ϵ the identity of G , and denote by $S(G)$ the set of subgroups of G . The subgroup generated by $g_1, \dots, g_k \in G$ is denoted by $\langle g_1, \dots, g_k \rangle$. We write $H \leq G$ if H is a subgroup of G , and write $H < G$ if H is a proper subgroup of G . The Möbius function on the poset $(S(G), \leq)$ of all subgroups of G is denoted by μ (see, e.g., [1]). We denote by $C(G)$ a full set of non-conjugate subgroups of G . Let $H \leq G$. We set ${}^gH = gHg^{-1}$ and $H^g = g^{-1}Hg$ for $g \in G$, and denote by (H) the set of conjugates of H in G . The normalizer of H in G is denoted by $N_G(H)$. We denote by $|G : H|$ the index of H in G , and denote by G/H the set of left cosets gH , $g \in G$, of H in G . Given $K, U \leq G$, $K \backslash G/U$ denotes the set of (K, U) -double cosets KgU , $g \in G$, in G . The category of finite left G -sets and G -equivariant maps is denoted by $G\text{-set}$. For each finite set X , we denote by $|X|$ the cardinality of X . The natural numbers, the rational integers, the rational numbers, and the complex numbers are denoted by $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$, and $\mathbb{C}$, respectively. We set $[n] = \{1, 2, \dots, n\}$ for each $n \in \mathbb{N}$. The identity map on a set Σ is denoted by id_Σ . For each group V , we denote by $\text{Hom}(V, \langle -1 \rangle)$ the group consisting of all group homomorphisms from V to the unit group $\langle -1 \rangle$ of $\mathbb{Z}$ with pointwise product.

2 Monomial Burnside rings

2A 1-cocycles

Throughout the paper, let G be a finite group, and let A be a finite G -group, that is, A is a finite group on which G acts via a homomorphism from G to the group of automorphisms of A (cf. [26, Chapter 1, Definition 8.1]). We start with the definition of (G, A) -sets introduced by Dress [12] (see also [27]). Given $g \in G$

and $a \in A$, the effect of g on a is denoted by ga . A finite free right A -set Y is called a (G, A) -set if it is also a left G -set and if

$$g(ya) = (gy) {}^ga$$

for all $g \in G$, $a \in A$, and $y \in Y$. A map between (G, A) -sets is called a (G, A) -equivariant map if it is a morphism of both left G -sets and right A -sets. We now obtain the category of (G, A) -sets such that the empty set is an initial object, which is denoted by (G, A) -**set**. Under the assumption that A is abelian, the set of isomorphism classes of (G, A) -sets forms a commutative unital semiring, and the monomial Burnside ring $\Omega(G, A)$ is defined to be the associated Grothendieck ring (cf. [12]).

For a (G, A) -set Y , we denote by Y/A the set of A -orbits $yA := \{ya \mid a \in A\}$, $y \in Y$, on Y , which is considered as a left G -set with the action of G given by

$$g(yA) = gyA$$

for all $g \in G$ and $y \in Y$. A (G, A) -set Y is said to be simple if Y/A is a transitive left G -set. Given a pair of (G, A) -sets Y_1 and Y_2 , their disjoint union $Y_1 \dot{\cup} Y_2$ is also a (G, A) -set. Every (G, A) -set is a disjoint union of simple (G, A) -sets. A subset of a (G, A) -set is said to be a (G, A) -subset if it is closed under the actions of G and A .

Let A° be the opposite group of A . For each $a \in A$, let a° denote the element of A° corresponding to a . By definition, $a^\circ b^\circ = (ba)^\circ$ for all $a, b \in A$. We view A° as a G -group with the action given by that of G on A , and denote by F the semidirect product $A^\circ \rtimes G$ of A° and G . Each (G, A) -set Y is viewed as a left F -set with the action of F given by

$$(a^\circ, g)y = (gy)a \quad (2.1)$$

for all $(a^\circ, g) \in F$ and $y \in Y$. A (G, A) -set is simple if and only if it is a transitive left F -set. A bijection between (G, A) -sets is an isomorphism of (G, A) -sets if and only if it is an isomorphism of left F -sets.

Let $H \leq G$. By restriction of operators from G to H , we view A as an H -group. A map $\sigma : H \rightarrow A$ is called a 1-cocycle or a crossed homomorphism if

$$\sigma(h_1 h_2) = \sigma(h_1) {}^{h_1}\sigma(h_2)$$

for all $h_1, h_2 \in H$ (cf. [26, I, p. 243]). We define a 1-cocycle $1_H : H \rightarrow A$ by $1_H(h) = \epsilon_A$ for all $h \in H$, where ϵ_A is the identity of A .

Definition 2.1 For each $H \leq G$, we denote by $Z^1(H, A)$ the set of 1-cocycles from H to A . Let $\mathcal{S}(G, A)$ be the set of pairs (H, σ) of $H \leq G$ and $\sigma \in Z^1(H, A)$. Given $(H, \sigma) \in \mathcal{S}(G, A)$, we fix a complete set $\{g_1, g_2, \dots, g_n\}$ with $g_1 = \epsilon$ of representatives of G/H , and define a (G, A) -set $(G/H)_\sigma$ to be the cartesian product $A \times (G/H)$ with the left action of G and the right action of A given by

$$g(a, g_j H) = ({}^{g_j'}\sigma(g_j'^{-1} g g_j) {}^ga, g_j' H) \quad \text{and} \quad (a, g_j H)b = (ab, g_j H),$$

where $gg_jH = g_{j'}H$, for all $g \in G$, $a, b \in A$, and $j \in [n]$, respectively.

Let $(H, \sigma) \in \mathcal{S}(G, A)$. Then $(G/H)_\sigma$ is a transitive left F -set. We define

$$F_{(H, \sigma)} := \{(\sigma(h)^{\circ-1}, h) \in F \mid h \in H\},$$

so that $F_{(H, \sigma)}$ is the stabilizer of $(\epsilon_A, H) \in (G/H)_\sigma$ in F (see [27, §2]), and make the set $F/F_{(H, \sigma)}$ of left cosets of $F_{(H, \sigma)}$ in F into a (G, A) -set by defining

$$g((a^\circ, r)F_{(H, \sigma)}) = ({}^g a^\circ, gr)F_{(H, \sigma)} \quad \text{and} \quad ((a^\circ, r)F_{(H, \sigma)})b = ((ab)^\circ, r)F_{(H, \sigma)} \quad (2.2)$$

for all $g \in G$, $b \in A$, and $(a^\circ, r) \in F$.

Lemma 2.2 *Let $(H, \sigma) \in \mathcal{S}(G, A)$. Then $(G/H)_\sigma \simeq F/F_{(H, \sigma)}$ as (G, A) -sets. In particular, the isomorphism class of (G, A) -sets containing $(G/H)_\sigma$ is independent of the choice of $g_2, \dots, g_n$ in Definition 2.1.*

Proof. There exists an isomorphism $F/F_{(H, \sigma)} \xrightarrow{\sim} (G/H)_\sigma$ of F -sets given by

$$(a^\circ, g)F_{(H, \sigma)} \mapsto (g(\epsilon_A, H))a$$

for all $(a^\circ, g) \in F$, because $F_{(H, \sigma)}$ is the stabilizer of $(\epsilon_A, H) \in (G/H)_\sigma$ in F . Thus we have $(G/H)_\sigma \simeq F/F_{(H, \sigma)}$ as (G, A) -sets, completing the proof. $\square$

Remark 2.3 Given a simple (G, A) -set Y and $y \in Y$, the stabilizer F_y of y in F coincides with $F_{(H, \sigma)}$ for some $(H, \sigma) \in \mathcal{S}(G, A)$ (see the proof of [27, Lemma 2.1]), and hence $Y \simeq F/F_{(H, \sigma)}$ as (G, A) -sets. Under the notation of Definition 2.1, we may define $(G/H)_\sigma$ without assuming that $g_1 = \epsilon$. In such a case, $F_{(H, \sigma)}$ is the stabilizer of $(\sigma(g_1)^{-1}, H) \in (G/H)_\sigma$ in F , which yields $(G/H)_\sigma \simeq F/F_{(H, \sigma)}$.

2B Isomorphism classes

We give a complete set of representatives of isomorphism classes of (G, A) -sets.

Definition 2.4 Let $(H, \sigma) \in \mathcal{S}(G, A)$. Suppose that $g \in G$ and $a \in A$. We define two 1-cocycles $g\sigma : {}^gH \rightarrow A$ and $\sigma^a : H \rightarrow A$ by

$$(g\sigma)(ghg^{-1}) = {}^g\sigma(h) \quad \text{and} \quad \sigma^a(h) = a^{-1}\sigma(h)h_a$$

for all $h \in H$, respectively.

Let $H \leq G$, and let $\sigma, \tau \in Z^1(H, A)$. We write $\sigma =_A \tau$ if $\tau = \sigma^a$ for some $a \in A$.

Lemma 2.5 *Let $(H, \sigma) \in \mathcal{S}(G, A)$. Then $g\sigma =_A g(\sigma^a)$ for any $g \in G$ and $a \in A$.*

Proof. We have $g(\sigma^a) = (g\sigma)^{ga}$ for any $g \in G$ and $a \in A$, completing the proof. $\square$

The argument of the proof of Lemma 2.5 ensures that $\mathcal{S}(G, A)$ is a left F -set with the action of F given by

$$(a^\circ, g)(H, \sigma) = ({}^gH, (g\sigma)^a)$$

for all $(a^\circ, g) \in F$ and $(H, \sigma) \in \mathcal{S}(G, A)$.

By [27, Lemma 2.3], (H, σ) and (U, τ) are contained in the same F -orbit on $\mathcal{S}(G, A)$ if and only if $(G/H)_\sigma \simeq (G/U)_\tau$ as (G, A) -sets.

Lemma 2.6 *Let $H \leq G$, and let $\sigma \in Z^1(H, A)$. Then $h\sigma = \sigma^{\sigma(h)}$ for any $h \in H$. Moreover, given $\sigma_0 \in Z^1(H, A)$, $\sigma_0 =_A \sigma$ if and only if $(H/H)_{\sigma_0} \simeq (H/H)_\sigma$.*

Proof. The first assertion is shown in the proof of [27, Lemma 3.2]. Suppose that $\sigma_0 \in Z^1(H, A)$. By [27, Lemma 2.3], $(H/H)_{\sigma_0} \simeq (H/H)_\sigma$ if and only if there exist some $h \in H$ and $a \in A$ such that $\sigma_0 = (h\sigma)^a$. Hence the second assertion follows from the first one. This completes the proof. $\square$

Definition 2.7 We define a subset $\mathcal{R}(G, A)$ of $\mathcal{S}(G, A)$ to be a complete set of representatives of F -orbits on $\mathcal{S}(G, A)$ such that $H \in \mathcal{C}(G)$ for any $(H, \sigma) \in \mathcal{R}(G, A)$.

The following proposition is [27, Proposition 2.4].

Proposition 2.8 *Let Y be a simple (G, A) -set. There exists a unique element (H, σ) of $\mathcal{R}(G, A)$ such that $Y \simeq (G/H)_\sigma$ as (G, A) -sets.*

Let $H \leq G$, and let $X \in H\text{-set}$. We define a left action of H on the cartesian product $G \times X$ of G and X by

$$h(g, x) = (gh^{-1}, hx)$$

for all $h \in H$ and $(g, x) \in G \times X$. Given $(g, x) \in G \times X$, let $g \otimes x$ denote the H -orbit containing (g, x) . The left G -set $\text{ind}_H^G(X)$ induced from X is the set of H -orbits on $G \times X$ with the action of G given by

$$g(r \otimes x) = gr \otimes x$$

for all $g, r \in G$ and $x \in X$ (cf. [11, §4]). Let $g \in G$, and set $g \otimes X = \{g \otimes x \mid x \in X\}$, which is a subset of $\text{ind}_H^G(X)$. The left gH -set $\text{con}_H^g(X)$ conjugate to X is the set $g \otimes X$ with the action of gH given by

$$ghg^{-1}(g \otimes x) = g \otimes hx$$

for all $h \in H$ and $x \in X$, and is denoted simply by gX .

Definition 2.9 Let $H \leq G$, and let T be an (H, A) -set. The (G, A) -set $\text{ind}_H^G(T)$ induced from T is the left G -set $\text{ind}_H^G(T)$ with the right action of A given by

$$(r \otimes t)a = r \otimes t^{r^{-1}}a$$

for all $r \in G$, $t \in T$, and $a \in A$ (cf. [27, Remark 6.2]). Let $g \in G$. The $({}^gH, A)$ -set $\text{con}_H^g(T)$ conjugate to T is the left gH -set gT with the right action of A given by

$$(g \otimes t)a = g \otimes t^{g^{-1}}a$$

for all $t \in T$ and $a \in A$ (cf. [27, Remark 6.4]), and is denoted simply by gT .

Lemma 2.10 If $U \leq H \leq G$ and $\tau \in Z^1(U, A)$, then $\text{ind}_H^G((H/U)_\tau) \simeq (G/U)_\tau$, ${}^g((H/U)_\tau) \simeq ({}^gH/{}^gU)_{g\tau}$ for each $g \in G$, and ${}^h((H/U)_\tau) \simeq (H/U)_\tau$ for all $h \in H$.

Proof. The proof is straightforward. Note that the last assertion follows from the second one and [27, Lemma 2.3]. $\square$

Let $(H, \sigma) \in \mathcal{S}(G, A)$, and let T be an (H, A) -set. For each $K \leq H$, we define a 1-cocycle $\sigma|_K : K \rightarrow A$, the restriction of σ , to be the map obtained by restriction of $\sigma : H \rightarrow A$ from H to K , and define a (K, A) -set $\text{res}_K^H(T)$, the restriction of T , to be the (K, A) -set T obtained by restriction of operators from H to K .

We show a Mackey decomposition formula for (G, A) -sets (cf. [27, Lemma 6.5]).

Lemma 2.11 Let $H \leq G$, and let $(U, \tau) \in \mathcal{S}(G, A)$. Then

$$\text{res}_H^G((G/U)_\tau) \simeq \dot{\bigcup}_{HgU \in H \backslash G/U} (H/(H \cap {}^gU))_{(g\tau)|_{H \cap {}^gU}},$$

where the disjoint union is taken over all (H, U) -double cosets HgU , $g \in G$, in G .

Proof. Let $\{g_1, g_2, \dots, g_m\}$ be a complete set of representatives of $H \backslash G/U$. For each $i \in [m]$, let $\{h_{i1}, h_{i2}, \dots, h_{i\ell_i}\}$ be a complete set of representatives of $H/(H \cap {}^{g_i}U)$. Then $\{h_{ij}g_i \mid i \in [m] \text{ and } j \in [\ell_i]\}$ is a complete set of representatives of G/U . We define a map $\Gamma : \text{res}_H^G((G/U)_\tau) \rightarrow \dot{\bigcup}_{i \in [m]} (H/(H \cap {}^{g_i}U))_{(g_i\tau)|_{H \cap {}^{g_i}U}}$ by

$$(a, h_{ij}g_iU) \mapsto (a, h_{ij}(H \cap {}^{g_i}U))$$

for all $i \in [m]$, $j \in [\ell_i]$, and $a \in A$. Obviously, this map is bijective and A -equivariant. Given $h \in H$, $i \in [m]$, and $j \in [\ell_i]$, if $hh_{ij} = h_{ij'}h' \in h_{ij'}(H \cap {}^{g_i}U)$ with $h' \in H \cap {}^{g_i}U$, then we have $h(h_{ij}g_i) = h_{ij'}g_i(g_i^{-1}h'g_i) \in h_{ij'}g_iU$ and

$${}^{h_{ij'}g_i}\tau((h_{ij}g_i)^{-1}h(h_{ij}g_i)) = {}^{h_{ij'}g_i}\tau(g_i^{-1}h'g_i) = {}^{h_{ij'}}(g_i\tau)(h_{ij'}^{-1}hh_{ij}).$$

Thus Γ is H -equivariant. (See also Lemma 2.2.) This completes the proof. $\square$

2C Tensor product

From now on, we assume that A is abelian. Hence $A = A^\circ$. Following [12], we define the tensor product $Y_1 \otimes Y_2$ of (G, A) -sets Y_1 and Y_2 . The cartesian product $Y_1 \times Y_2$ is viewed as a free right A -set with the action of A given by

$$(y_1, y_2)a = (y_1a^{-1}, y_2a)$$

for all $a \in A$ and $(y_1, y_2) \in Y_1 \times Y_2$. For each $(y_1, y_2) \in Y_1 \times Y_2$, let $y_1 \otimes y_2$ be the A -orbit containing (y_1, y_2) . We set

$$Y_1 \otimes Y_2 = \{y_1 \otimes y_2 \mid (y_1, y_2) \in Y_1 \times Y_2\},$$

and make it into a (G, A) -set by defining

$$g(y_1 \otimes y_2) = gy_1 \otimes gy_2 \quad \text{and} \quad (y_1 \otimes y_2)a = y_1 \otimes y_2a$$

for all $g \in G$, $a \in A$, and $(y_1, y_2) \in Y_1 \times Y_2$. These actions are well-defined, because

$$g((y_1b^{-1} \otimes y_2b)a) = g(y_1b^{-1}) \otimes g(y_2ba) = (gy_1)^{gb^{-1}} \otimes g(y_2a)^{gb} = g((y_1 \otimes y_2)a)$$

for all $g \in G$, $a, b \in A$, and $(y_1, y_2) \in Y_1 \times Y_2$. Obviously, $Y_1 \otimes Y_2 \simeq Y_2 \otimes Y_1$.

Lemma 2.12 *Let $K \leq H \leq G$, and let $g \in G$. For any (H, A) -sets T_1 and T_2 ,*

$$\text{res}_K^H(T_1 \otimes T_2) \simeq \text{res}_K^H(T_1) \otimes \text{res}_K^H(T_2) \quad \text{and} \quad {}^g(T_1 \otimes T_2) \simeq {}^gT_1 \otimes {}^gT_2.$$

Proof. The proof is straightforward. $\square$

Let $\mathbf{F}(G, A)$ be the free abelian group on the set of isomorphism classes of (G, A) -sets. For each (G, A) -set Y , we denote by $\overline{Y}$ the isomorphism class of (G, A) -sets containing Y . Let $\mathbf{F}(G, A)_0$ be the subgroup of $\mathbf{F}(G, A)$ generated by the elements

$$\overline{Y_1 \dot{\cup} Y_2} - \overline{Y_1} - \overline{Y_2}$$

for (G, A) -sets Y_1 and Y_2 . We define multiplication on the generators of $\mathbf{F}(G, A)$ by

$$\overline{Y_1} \cdot \overline{Y_2} = \overline{Y_1 \otimes Y_2}$$

for all (G, A) -sets Y_1 and Y_2 , and extend it to $\mathbf{F}(G, A)$ by linearity. Then $\mathbf{F}(G, A)$ is a commutative unital ring; moreover, $\mathbf{F}(G, A)_0$ is an ideal of $\mathbf{F}(G, A)$.

Definition 2.13 We define a commutative unital ring $\Omega(G, A)$ to be the quotient $\mathbf{F}(G, A)/\mathbf{F}(G, A)_0$, which is the ring of monomial representations of G with coefficients in A introduced by Dress [12] (see also [2]).

When $A = \{\epsilon_A\}$, which is the group consisting of only the identity, $\Omega(G, A)$ is isomorphic to the Burnside ring $\Omega(G)$ (see §5C).

For each (G, A) -set Y , we denote by $[Y]$ the coset $\overline{Y} + \mathbf{F}(G, A)_0$ of $\mathbf{F}(G, A)_0$ in $\mathbf{F}(G, A)$. By [12, Proposition 1(b)] (or [27, Lemma 2.6]), $[Y_1] = [Y_2]$ if and only if $\overline{Y_1} = \overline{Y_2}$. Multiplication on the generators of $\Omega(G, A)$ is given by

$$[Y_1] \cdot [Y_2] = [Y_1 \otimes Y_2]$$

for all (G, A) -sets Y_1 and Y_2 . The identity of $\Omega(G, A)$ is $[(G/G)_{1_G}]$.

A $\mathbb{Z}$ -lattice is a finitely generated $\mathbb{Z}$ -free $\mathbb{Z}$ -module. Obviously, $\Omega(G, A)$ is a $\mathbb{Z}$ -lattice. The statement of the following proposition is given in [12, Proposition 1(a)] (see also [2, Remark 2.2] and [27, Proposition 2.7]).

Proposition 2.14 *The elements $[(G/H)_\sigma]$ for $(H, \sigma) \in \mathcal{R}(G, A)$ form a free $\mathbb{Z}$ -basis of the $\mathbb{Z}$ -lattice $\Omega(G, A)$.*

Proof. The assertion follows from Proposition 2.8. $\square$

We obtain a product formula of simple (G, A) -sets (see also [2, Remark 2.3]).

Lemma 2.15 *Let $(H, \sigma), (U, \tau) \in \mathcal{S}(G, A)$. Then*

$$(G/H)_\sigma \otimes (G/U)_\tau \simeq \bigcup_{HgU \in H \backslash G/U} (G/(H \cap {}^gU))_{\sigma \cdot (g\tau)},$$

where $\sigma \cdot (g\tau) : H \cap {}^gU \rightarrow A$ is the pointwise product of $\sigma|_{H \cap {}^gU}$ and $(g\tau)|_{H \cap {}^gU}$.

Proof. We view the tensor product $(F/F_{(H,\sigma)}) \otimes (F/F_{(U,\tau)})$ of (G, A) -sets $F/F_{(H,\sigma)}$ and $F/F_{(U,\tau)}$ as a left F -set. The left F -set $(F/F_{(H,\sigma)}) \otimes (F/F_{(U,\tau)})$ is expressed as a disjoint union of F -orbits. We identify each $g \in G$ with $(\epsilon_A, g) \in F$ for shortness' sake. For any $(a, g), (b, r) \in F$,

$$(a, g)^{-1}((a, g)F_{(H,\sigma)} \otimes (b, r)F_{(U,\tau)}) = ({}^g{}^{-1}b, \epsilon)(F_{(H,\sigma)} \otimes g^{-1}rF_{(U,\tau)})$$

(see Eqs. (2.1) and (2.2)), which means that there exists an F -orbit containing both $(a, g)F_{(H,\sigma)} \otimes (b, r)F_{(U,\tau)}$ and $F_{(H,\sigma)} \otimes g^{-1}rF_{(U,\tau)}$. Let $g, r \in G$. Suppose that

$$F_{(H,\sigma)} \otimes rF_{(U,\tau)} = (a, h)(F_{(H,\sigma)} \otimes gF_{(U,\tau)}) = hF_{(H,\sigma)} \otimes (a, hg)F_{(U,\tau)}$$

with $(a, h) \in F$. Then $h \in H$ and $r^{-1}hg \in U$, which yields $g \in HrU$. Conversely, if $g \in HrU$ and $r^{-1}hg \in U$ with $h \in H$, then we have

$$F_{(H,\sigma)} \otimes rF_{(U,\tau)} = ({}^{hg}\tau(g^{-1}h^{-1}r)\sigma(h)^{-1}, h)(F_{(H,\sigma)} \otimes gF_{(U,\tau)}).$$

Consequently, both $F_{(H,\sigma)} \otimes rF_{(U,\tau)}$ and $F_{(H,\sigma)} \otimes gF_{(U,\tau)}$ are contained in the same F -orbit if and only if $g \in HrU$. Suppose that

$$F_{(H,\sigma)} \otimes gF_{(U,\tau)} = (a, h)(F_{(H,\sigma)} \otimes gF_{(U,\tau)}) = (a, h)F_{(H,\sigma)} \otimes hgF_{(U,\tau)}$$

with $(a, h) \in F$. Then there exists some $b \in A$ such that

$$(F_{(H,\sigma)}, gF_{(U,\tau)}) = (((a, h)F_{(H,\sigma)})b^{-1}, (hgF_{(U,\tau)})b) = ((b^{-1}a, h)F_{(H,\sigma)}, (b, hg)F_{(U,\tau)}),$$

which yields $h \in H \cap {}^gU$ and

$$(\sigma \cdot (g\tau))(h) = \sigma(h) {}^g\tau(g^{-1}hg) = (a^{-1}b) {}^g(g^{-1}b^{-1}) = a^{-1}.$$

Hence $(a, h) \in F_{(H \cap {}^gU, \sigma \cdot (g\tau))}$. Moreover, it is easily verified that $F_{(H \cap {}^gU, \sigma \cdot (g\tau))}$ is the stabilizer of $F_{(H,\sigma)} \otimes gF_{(U,\tau)}$. Thus it turns out that

$$(F/F_{(H,\sigma)}) \otimes (F/F_{(U,\tau)}) \simeq \bigcup_{HgU \in H \backslash G/U} F/F_{(H \cap {}^gU, \sigma \cdot (g\tau))}$$

as left F -sets. The lemma now follows from Lemma 2.2. This completes the proof. $\square$

For each $K \leq H \leq G$ and $g \in G$, there are additive maps

$$\begin{aligned} \text{con}_H^g : \Omega(H, A) &\rightarrow \Omega({}^gH, A), & \sum_T \ell_T[T] &\mapsto \sum_T \ell_T[\text{con}_H^g(T)], \\ \text{res}_K^H : \Omega(H, A) &\rightarrow \Omega(K, A), & \sum_T \ell_T[T] &\mapsto \sum_T \ell_T[\text{res}_K^H(T)], \quad \text{and} \\ \text{ind}_K^H : \Omega(K, A) &\rightarrow \Omega(H, A), & \sum_S k_S[S] &\mapsto \sum_S k_S[\text{ind}_K^H(S)], \end{aligned}$$

where $S \in (K, A)\text{-set}$, $T \in (H, A)\text{-set}$, and $k_S, \ell_T \in \mathbb{Z}$; these maps are called the conjugation map, the restriction map, and the induction map, respectively. By Lemma 2.12, conjugation maps and restriction maps are ring homomorphisms.

Proposition 2.16 *The family of $\mathbb{Z}$ -algebras $\Omega(H, A)$ for $H \leq G$, together with conjugation, restriction, and induction maps, defines a Green functor on G .*

Proof. The axioms of Green functor follow from Lemmas 2.10, 2.11, and 2.15 (cf. [4, 1.1. Definition]). As for the Frobenius axiom, we have

$$\text{res}_K^G((G/H)_\sigma \otimes (K/U)_\tau) \simeq \bigcup_{KgH \in K \backslash G/H} \bigcup_{L_g eU \in L_g \backslash K/U} (K/({}^gH \cap {}^eU))_{(g\sigma)|_{L_g} \cdot (e\tau)},$$

where $L_g = K \cap {}^gH$, and

$$\text{ind}_K^G(\text{res}_K^G((G/H)_\sigma \otimes (K/U)_\tau)) \simeq (G/H)_\sigma \otimes \text{ind}_K^G((K/U)_\tau)$$

for all $K \leq G$, $(H, \sigma) \in \mathcal{S}(G, A)$, and $(U, \tau) \in \mathcal{S}(K, A)$, completing the proof. $\square$

3 Multiplicative induction

3A Tensor induction

To begin with, we review the multiplicative induction $\text{Jnd}_H^G : X \mapsto \text{Map}_H(G, X)$, where $H \leq G$ and $X \in H\text{-set}$, given in [32, §3(a.3)] (see also [11, §4]).

Definition 3.1 Let $H \leq G$, and let T be an (H, A) -set. We define a left G -set $\text{Map}_H(G, T)$ to be the set of maps $f : G \rightarrow T$ such that $f(hg) = hf(g)$ for all $h \in H$ and $g \in G$ with the action of G given by

$$(gf)(r) = f(rg)$$

for all $g, r \in G$ and $f \in \text{Map}_H(G, T)$.

Remark 3.2 Under the notation of Definition 3.1, the left G -set $\text{Map}_H(G, T)$ is viewed as a (G, A) -set with the right action of A given by

$$(fa)(r) = f(r)^{ra}$$

for all $r \in G$, $a \in A$, and $f \in \text{Map}_H(G, T)$. However, we need hardly recall such a right action of A on $\text{Map}_H(G, T)$ (see Definitions 3.3 and 3.5) in relation to multiplicative induction for monomial Burnside rings (see Proposition 3.20).

Let $H \leq G$, and let T be an (H, A) -set. The tensor induced G -set $T^{\otimes G}$ obtained from T (see [8, §80C]) is isomorphic to $\text{Map}_H(G, T)$ and is related to tensor induction of modules. By modifying $\text{Map}_H(G, T)$, we define tensor induction for (H, A) -sets, and then define multiplicative induction for monomial Burnside rings in §3C.

Let Hg , $g \in G$, be the right coset of H in G containing g . Given $g, r \in G$ with $Hg \neq Hr$ and $a \in A$, we define a relation $\sim_{(g,r,a)}$ on $\text{Map}_H(G, T)$ by

$$\begin{aligned} f \sim_{(g,r,a)} f' : \iff & \quad f(hg)^{ha} = f'(hg) \text{ and } f(hr) = f'(hr)^{hr}a \text{ for all } h \in H, \\ & \text{and } f(g') = f'(g') \text{ for all } g' \in G - Hg \dot{\cup} Hr. \end{aligned}$$

Let $\sim_A$ be the equivalence relation on $\text{Map}_H(G, T)$ generated by the relations $\sim_{(g,r,a)}$ for $g, r \in G$ and $a \in A$. For each $f \in \text{Map}_H(G, T)$, we denote by $\hat{f}$ the equivalence class containing f with respect to the equivalence relation $\sim_A$.

Definition 3.3 Let $H \leq G$, and let T be an (H, A) -set. We define

$$\widehat{\text{Map}}_H(G, T) := \{\hat{f} \mid f \in \text{Map}_H(G, T)\},$$

and make it into a free right A -set by defining

$$\hat{f}a = \hat{f}_a \quad \text{with} \quad f_a : G \rightarrow T, \quad r \mapsto f_a(r) = \begin{cases} f(r)^{ra} & \text{if } r \in H, \\ f(r) & \text{if } r \in G - H \end{cases} \quad (3.1)$$

for all $a \in A$ and $f \in \text{Map}_H(G, T)$.

The following lemma tells us of a suitable left action of G on the free right A -set $\widehat{\text{Map}}_H(G, T)$ defined above for an extension of the multiplicative induction $\text{Jnd}_H^G : X \mapsto \text{Map}_H(G, X)$ where $X \in H\text{-set}$.

Lemma 3.4 *Let $H \leq G$, and let T be an (H, A) -set. Then $\widehat{gf}_a = \widehat{gf}^a$, where f_a is given in Eq.(3.1), for all $g \in G$, $a \in A$, and $f \in \text{Map}_H(G, T)$.*

Proof. Suppose that $g \in G$, $a \in A$, and $f \in \text{Map}_H(G, T)$. By definition,

$$(gf_a)(r) = \begin{cases} f(rg)^{r^g a} & \text{if } rg \in H, \\ f(rg) & \text{if } rg \in G - H, \end{cases}$$

and

$$(gf)^a(r) = \begin{cases} f(rg)^{r^g a} & \text{if } r \in H, \\ f(rg) & \text{if } r \in G - H. \end{cases}$$

Hence we may assume that $g \notin H$. Observe that for any $h \in H$,

$$(gf_a)(h)^h({}^g a) = (gf)^a(h) \quad \text{and} \quad (gf_a)(hg^{-1}) = (gf)^a(hg^{-1})^{hg^{-1}({}^g a)}.$$

Moreover, $(gf_a)(r) = (gf)^a(r)$ for all $r \in G - H \cup Hg^{-1}$. Thus $gf_a \sim_{(\epsilon, g^{-1}, {}^g a)} (gf)^a$. We now obtain $\widehat{gf}_a = \widehat{gf}^a$, completing the proof. $\square$

Definition 3.5 (Tensor induction) Let $H \leq G$, and let T be an (H, A) -set. We make the free right A -set $\widehat{\text{Map}}_H(G, T)$ into a left G -set by defining

$$g\widehat{f} = \widehat{gf}$$

for all $g \in G$ and $f \in \text{Map}_H(G, T)$, so that $\widehat{\text{Map}}_H(G, T)$ is a (G, A) -set. The operation which assigns to T the (G, A) -set $\widehat{\text{Map}}_H(G, T)$ is called tensor induction (cf. [8, §80C]), and is related to tensor induction for 1-cocycles (see §3B).

Remark 3.6 Keep the notation of Definition 3.5, and assume further that G acts trivially on A . Then the (G, A) -sets are considered as the A -fibred G -sets defined by Barker [2, §2], and the (G, A) -set $\widehat{\text{Map}}_H(G, T)$ obtained from T by tensor induction is identified with the A -fibred G -set $\text{Ten}_H^G(T)$ defined by Barker [2, §9].

We present a fundamental lemma which is essential to the investigation of multiplicative induction for monomial Burnside rings.

Lemma 3.7 *Let $H \leq G$, and let T be an (H, A) -set. Suppose that $\{g_1, g_2, \dots, g_n\}$ with $g_1 = \epsilon$ is a complete set of representatives of G/H . Let $f \in \text{Map}_H(G, T)$, and define $f^{(0)} \in \text{Map}_H(G, T)$ by $f^{(0)}(hg_j^{-1}) = f(hg_j^{-1})^{h a_j}$ with $a_j \in A$ for all $h \in H$ and $j \in [n]$. Then $f^{(0)} \sim_A f_a$, where $a = {}^{g_1}a_1 {}^{g_2}a_2 \dots {}^{g_n}a_n$, and hence $\widehat{f^{(0)}} = \widehat{f}a$.*

Proof. For each integer k with $1 \leq k \leq n$, we define $f^{(k)} \in \text{Map}_H(G, T)$ by

$$f^{(k)}(hg_j^{-1}) = \begin{cases} f(hg_j^{-1}) & \text{if } j \in [k], \\ f^{(0)}(hg_j^{-1}) & \text{if } j = k+1, k+2, \dots, n \end{cases}$$

for all $h \in H$. In particular, $f^{(n)} = f$. Obviously, $f^{(1)} = f_{g_1 a_1^{-1}}^{(0)}$. Let k be an integer with $2 \leq k \leq n$. Then

$$f^{(k)}(hg_k^{-1}) = f^{(k-1)}(hg_k^{-1}) h a_k^{-1} \quad \text{and} \quad f^{(k)}(hg_j^{-1}) = f^{(k-1)}(hg_j^{-1})$$

for all $h \in H$ and $j \in [n]$ with $j \neq k$, and

$$f_{g_k a_k^{-1}}^{(k-1)}(hg_1^{-1}) = f^{(k-1)}(hg_1^{-1}) h g_k a_k^{-1} \quad \text{and} \quad f_{g_k a_k^{-1}}^{(k-1)}(hg_j^{-1}) = f^{(k-1)}(hg_j^{-1})$$

for all $h \in H$ and $j = 2, 3, \dots, n$. This shows that $f_{g_k a_k^{-1}}^{(k-1)} \sim_{(g_k^{-1}, g_1^{-1}, g_k a_k^{-1})} f^{(k)}$. Hence we have $f^{(0)} \sim_A f_a$, completing the proof. $\square$

Remark 3.8 Let $H \leq G$, and let T be an (H, A) -set. By Lemma 3.7, we have

$$|\widehat{\text{Map}}_H(G, T)/A| = |\text{Map}_H(G, T/A)|,$$

whence

$$|\widehat{\text{Map}}_H(G, T)| = |T/A|^{[G/H]} \cdot |A|.$$

The following proposition, which is a generalization of [32, §3(a.13)], describes a Mackey decomposition formula (see also [2, Lemma 9.1]).

Proposition 3.9 *Let $H, K \leq G$. For each (H, A) -set T ,*

$$\text{res}_K^G(\widehat{\text{Map}}_H(G, T)) \simeq \bigotimes_{KgH \in K \backslash G/H} \widehat{\text{Map}}_{K \cap {}^{g_i}H}(K, \text{res}_{K \cap {}^{g_i}H}^{{}^{g_i}H}({}^{g_i}T)).$$

Proof. Let $\{g_1, g_2, \dots, g_m\}$ with $g_1 = \epsilon$ be a complete set of representatives of $K \backslash G/H$. For each $i \in [m]$, let $\{r_{i1}, r_{i2}, \dots, r_{i\ell_i}\}$ be a complete set of representatives of $K/(K \cap {}^{g_i}H)$. Then $\{r_{ij}g_i \mid i \in [m] \text{ and } j \in [\ell_i]\}$ is a complete set of representatives of G/H . Let $i \in [m]$. There is a map

$$\Phi_i : \text{res}_K^G(\text{Map}_H(G, T)) \rightarrow \text{Map}_{K \cap {}^{g_i}H}(K, \text{res}_{K \cap {}^{g_i}H}^{{}^{g_i}H}({}^{g_i}T))$$

given by

$$\Phi_i(f)({}^{g_i}hr_{ij}^{-1}) = g_i \otimes f(h(r_{ij}g_i)^{-1}) (= {}^{g_i}h(g_i \otimes f((r_{ij}g_i)^{-1}))) \in {}^{g_i}T$$

for all $h \in {}^{g_i^{-1}}K \cap H$, $j \in [\ell_i]$, and $f \in \text{Map}_H(G, T)$. Given $j \in [\ell_i]$, we have

$$\Phi_i(rf)(r_{ij}^{-1}) = g_i \otimes f((r_{ij}g_i)^{-1}r) = g_i \otimes f(g_i^{-1}(r_{ij}^{-1}rr_{ij'})g_i(r_{ij'}g_i)^{-1})$$

and

$$(r\Phi_i(f))(r_{ij}^{-1}) = \Phi_i(f)(r_{ij}^{-1}r) = \Phi_i(f)((r_{ij}^{-1}rr_{ij'})r_{ij'}^{-1}),$$

where $r_{ij}(K \cap {}^{g_i}H) = rr_{ij'}(K \cap {}^{g_i}H)$, for all $r \in K$ and $f \in \text{Map}_H(G, T)$. Thus Φ_i is a K -equivariant map. We now define a K -equivariant map

$$\widehat{\Phi} : \text{res}_K^G(\widehat{\text{Map}}_H(G, T)) \rightarrow \bigotimes_{i=1}^m \widehat{\text{Map}}_{K \cap {}^{g_i}H}(K, \text{res}_{K \cap {}^{g_i}H}^{g_i H}({}^{g_i}T))$$

by

$$\widehat{f} \mapsto \widehat{\Phi_1(f)} \otimes \widehat{\Phi_2(f)} \otimes \cdots \otimes \widehat{\Phi_m(f)}$$

for all $f \in \text{Map}_H(G, T)$. (Of course this map is well-defined; see Remark 3.10.) The map $\widehat{\Phi}$ is also a (K, A) -equivariant map, because

$$(\widehat{\Phi_1(f)} \otimes \widehat{\Phi_2(f)} \otimes \cdots \otimes \widehat{\Phi_m(f)})a = \widehat{\Phi_1(f)_a} \otimes \widehat{\Phi_2(f)} \otimes \cdots \otimes \widehat{\Phi_m(f)}$$

and

$$\Phi_i(f_a)(r_{ij}^{-1}) = \begin{cases} \epsilon \otimes f(r_{1j}^{-1})r_{1j}^{-1}a & = \Phi_1(f)_a(r_{1j}^{-1}) & \text{if } i = 1 \text{ and } r_{1j} \in H, \\ \epsilon \otimes f(r_{1j}^{-1}) & = \Phi_1(f)_a(r_{1j}^{-1}) & \text{if } i = 1 \text{ and } r_{1j} \notin H, \\ g_i \otimes f((r_{ij}g_i)^{-1}) & = \Phi_i(f)(r_{ij}^{-1}) & \text{if } i \neq 1 \end{cases}$$

for all $i \in [m]$, $j \in [\ell_i]$, $a \in A$, and $f \in \text{Map}_H(G, T)$. Thus it only remains for us to show that $\widehat{\Phi}$ is bijective. For each $i \in [m]$, choose $f_i \in \text{Map}_{K \cap {}^{g_i}H}(K, \text{res}_{K \cap {}^{g_i}H}^{g_i H}({}^{g_i}T))$. Given $i \in [m]$ and $j \in [\ell_i]$, we suppose that $f_i(r_{ij}^{-1}) = g_i \otimes t_{ij} \in {}^{g_i}T$ with $t_{ij} \in T$. Now define $f \in \text{Map}_H(G, T)$ by

$$f(h(r_{ij}g_i)^{-1}) = ht_{ij} \in T$$

for all $h \in H$, $i \in [m]$, and $j \in [\ell_i]$. Then $\widehat{\Phi(f)} = \widehat{f_1} \otimes \widehat{f_2} \otimes \cdots \otimes \widehat{f_m}$. Thus $\widehat{\Phi}$ is surjective, which means that it is also injective, because

$$|\widehat{\text{Map}}_H(G, T)| = |T/A|^{\sum_{i=1}^m \ell_i} \cdot |A| = \left| \bigotimes_{i=1}^m \widehat{\text{Map}}_{K \cap {}^{g_i}H}(K, \text{res}_{K \cap {}^{g_i}H}^{g_i H}({}^{g_i}T)) \right|$$

by Remark 3.8. We now conclude that $\widehat{\Phi}$ is bijective, completing the proof. $\square$

Remark 3.10 In the proof of Proposition 3.9, assume that $f \sim_{((r_{ij}g_i)^{-1}, (r_{i'j'}g_{i'})^{-1}, a)} f'$ with $f, f' \in \text{Map}_H(G, T)$ and $a \in A$. Let $u \in [m]$ and $v \in [\ell_u]$. Then we have

$$\Phi_u(f)({}^{g_u}hr_{uv}^{-1}) = \begin{cases} \Phi_i(f')({}^{g_i}hr_{ij}^{-1}){}^{g_i}hr_{ij}^{-1}a^{-1} & \text{if } (u, v) = (i, j), \\ \Phi_{i'}(f')({}^{g_{i'}}hr_{i'j'}^{-1}){}^{g_{i'}}hr_{i'j'}^{-1}a & \text{if } (u, v) = (i', j'), \\ \Phi_u(f')({}^{g_u}hr_{uv}^{-1}) & \text{otherwise} \end{cases}$$

for all $h \in {}^{g_u^{-1}}K \cap H$. This, combined with Lemma 3.7, shows that

$$\widehat{\Phi_u(f)} = \widehat{\Phi_u(f')} \quad \text{if } u \neq i, i', \quad \widehat{\Phi_i(f)} = \widehat{\Phi_i(f')}a^{-1}, \quad \text{and} \quad \widehat{\Phi_{i'}(f)} = \widehat{\Phi_{i'}(f')}a.$$

Hence we have $\widehat{\Phi}(f) = \widehat{\Phi}(f')$. Consequently, the map $\widehat{\Phi}$ is well-defined.

3B Tensor induction for 1-cocycles

We introduce tensor induction for 1-cocycles, and see that it is closely allied to tensor induction for (H, A) -sets with $H \leq G$.

Definition 3.11 Let $(H, \sigma) \in \mathcal{S}(G, A)$. We fix a complete set $\{g_1, g_2, \dots, g_n\}$ with $g_1 = \epsilon$ of representatives of G/H , and define a 1-cocycle $\sigma^{\otimes G} : G \rightarrow A$ by

$$\sigma^{\otimes G}(g) = \prod_{j=1}^n {}^{g_j'}\sigma(g_j^{-1}gg_j),$$

where $gg_jH = g_j'H$, for all $g \in G$. The operation which assigns to σ the 1-cocycle $\sigma^{\otimes G} : G \rightarrow A$ is called tensor induction (cf. [8, §13A]).

Remark 3.12 Keep the notation of Definition 3.11, and let $h_1, h_2, \dots, h_n \in H$. Then

$$(\sigma^{\otimes G})^a(g) = \prod_{j=1}^n {}^{g_j'h_j'}\sigma(h_j^{-1}g_j'^{-1}gg_jh_j) \quad \text{with} \quad a = \prod_{j=1}^n {}^{g_j'}\sigma(h_j)$$

for all $g \in G$ (see Definition 2.4), because

$$\begin{aligned} {}^{g_j'h_j'}\sigma(h_j^{-1}g_j'^{-1}gg_jh_j) &= {}^{g_j'h_j'}\sigma(h_j'^{-1}) {}^{g_j'}\sigma(g_j'^{-1}gg_jh_j) \\ &= {}^{g_j'}\sigma(h_j')^{-1} {}^{g_j'}\sigma(g_j'^{-1}gg_j) {}^{gg_j}\sigma(h_j) \end{aligned}$$

for all $j \in [n]$. Hence the subset $\{(\sigma^{\otimes G})^a \mid a \in A\}$ of $Z^1(G, A)$ is independent of the choice of a complete set of representatives of G/H . Likewise, if $b \in A$, then

$$(\sigma^{\otimes G})^c = (\sigma^b)^{\otimes G} \quad \text{with} \quad c = \prod_{j=1}^n {}^{g_j'}b.$$

Example 3.13 Let $(H, \sigma) \in \mathcal{S}(G, A)$. Obviously, A is a free right A -set with the action given by the product operation on A . We make it into an (H, A) -set $A^{(\sigma)}$ isomorphic to $(H/H)_\sigma$ by defining

$$ha = \sigma(h) {}^h a$$

for all $h \in H$ and $a \in A^{(\sigma)}$. For any $K \leq H$, $\text{res}_K^H(A^{(\sigma)}) = A^{(\sigma|_K)}$ (see Lemma 2.11). Keep the notation of Definition 3.11, and identify $(H/H)_\sigma$ with $A^{(\sigma)}$. We define an element $\tilde{\sigma}$ of $\text{Map}_H(G, A^{(\sigma)})$ by

$$\tilde{\sigma}(hg_i^{-1}) = \sigma(h)$$

for all $h \in H$ and $i \in [n]$. Let $f \in \text{Map}_H(G, A^{(\sigma)})$. For each $j \in [n]$, we set $a_j = f(g_j^{-1}) \in A^{(\sigma)}$. Since $f(hg_j^{-1}) = \tilde{\sigma}(hg_j^{-1})^h a_j$ for all $h \in H$ and $j \in [n]$, it follows from Lemma 3.7 with $f^{(0)} = f$ that $\hat{f} = \hat{\sigma}a$ where $a = {}^{g_1}a_1 {}^{g_2}a_2 \cdots {}^{g_n}a_n$. Hence $\widehat{\text{Map}}_H(G, A^{(\sigma)}) = \{\hat{\sigma}a \mid a \in A\}$. Let $g \in G$. We have

$$(g\tilde{\sigma})(hg_j^{-1}) = \tilde{\sigma}(hg_j^{-1}g) = \sigma(hg_j^{-1}gg_{j'}) = \sigma(h)^h \sigma(g_j^{-1}gg_{j'}) = \tilde{\sigma}(hg_j^{-1})^h \sigma(g_j^{-1}gg_{j'}),$$

where $g_jH = gg_{j'}H$, for all $h \in H$ and $j \in [n]$. Thus it follows from Lemma 3.7 that $g\hat{\sigma} = \hat{\sigma}\sigma^{\otimes G}(g)$. Moreover, there exists an isomorphism $\widehat{\text{Map}}_H(G, A^{(\sigma)}) \xrightarrow{\sim} A^{(\sigma^{\otimes G})}$ of (G, A) -sets given by

$$\hat{\sigma}a \mapsto a$$

for all $a \in A$. Thus $\widehat{\text{Map}}_H(G, (H/H)_\sigma) \simeq (G/G)_{\sigma^{\otimes G}}$.

The following proposition describes a Mackey decomposition formula.

Proposition 3.14 *Let $H, K \leq G$. For each $\sigma \in Z^1(H, A)$,*

$$\sigma^{\otimes G}|_K =_A \prod_{KgH \in K \backslash G/H} (g\sigma)|_{K \cap {}^gH}^{\otimes K}.$$

Proof. By Lemma 2.10, Proposition 3.9, and Example 3.13, we have

$$(K/K)_{\sigma^{\otimes G}|_K} \simeq \bigotimes_{KgH \in K \backslash G/H} (K/K)_{(g\sigma)|_{K \cap {}^gH}}^{\otimes K},$$

which, combined with Lemma 2.15, implies that

$$(K/K)_{\sigma^{\otimes G}|_K} \simeq (K/K) \prod_{KgH \in K \backslash G/H} (g\sigma)|_{K \cap {}^gH}^{\otimes K}.$$

The assertion follows from this fact and Lemma 2.6. This completes the proof. $\square$

The following lemma states basic properties of tensor induction for 1-cocycles.

Lemma 3.15 *Let $U \leq K \leq H$, and let $g \in G$. Then*

$$g\nu^{\otimes H} =_A (g\nu)^{\otimes {}^gH} \quad \text{and} \quad (\tau^{\otimes K})^{\otimes H} =_A \tau^{\otimes H}$$

for all $\nu \in Z^1(K, A)$ and $\tau \in Z^1(U, A)$.

Proof. Fix a complete set $\{h_1, h_2, \dots, h_m\}$ with $h_1 = \epsilon$ of representatives of H/K and a complete set $\{r_1, r_2, \dots, r_k\}$ with $r_1 = \epsilon$ of representatives of K/U . Given $\nu \in Z^1(K, A)$ and $\tau \in Z^1(U, A)$, we have

$$\begin{aligned} (g\nu^{\otimes H})(g_h) &= \prod_{j=1}^m g_{h_{j'}}\nu(h_{j'}^{-1}hh_j) \\ &= \prod_{j=1}^m g_{h_{j'}}g_\nu((h_{j'}^{-1}g^{-1})g_h(g_hh_j)) \\ &= \prod_{j=1}^m g_{h_{j'}}g_\nu(g^{-1}(g_{h_{j'}}^{-1}g_hg_hh_j)g) \\ &= \prod_{j=1}^m g_{h_{j'}}(g\nu)(g_{h_{j'}}^{-1}g_hg_hh_j), \end{aligned}$$

where $hh_jK = h_{j'}K$, and

$$\begin{aligned} (\tau^{\otimes K})^{\otimes H}(h) &= \prod_{j=1}^m h_{j'}\tau^{\otimes K}(h_{j'}^{-1}hh_j) \\ &= \prod_{j=1}^m \prod_{i=1}^k h_{j'}r_{i'}\tau((h_{j'}r_{i'})^{-1}h(h_jr_i)), \end{aligned}$$

where $(h_{j'}^{-1}hh_j)r_iU = r_{i'}U$, for all $h \in H$. Consequently, the assertions follow from Remark 3.12. This completes the proof. $\square$

3C Algebraic maps

We define a subset $\Omega(G, A)^+$ of $\Omega(G, A)$ to be the set consisting of all elements $\sum_{(U, \tau) \in \mathcal{R}(G, A)} \ell_{(U, \tau)}[(G/U)_\tau]$ with $\ell_{(U, \tau)} \geq 0$, which is an additive semigroup. By Lemma 2.15, $\Omega(G, A)^+$ is closed under multiplication. For each $H \leq G$, there is a map (tensor induction) $\widehat{\text{Map}}_H(G, -) : \Omega(H, A)^+ \rightarrow \Omega(G, A)$ given by

$$[T] \mapsto [\widehat{\text{Map}}_H(G, T)]$$

for all (H, A) -sets T (cf. [8, (80.42)]). This map is multiplicative (see Lemma 3.19).

We review the concept of algebraic maps which is due to Dress [11]. Let B be an additive semigroup with zero element, and let E be an additive group. Given $c \in B$ and a map $f : B \rightarrow E$, we define a map $D_c f : B \rightarrow E$ by

$$d \mapsto f(c + d) - f(d)$$

for all $d \in B$. A map $f : B \rightarrow E$ is said to be algebraic of degree n if n is the least integer such that

$$D_{c_1} D_{c_2} \cdots D_{c_{n+1}} f = 0$$

for all $c_1, c_2, \dots, c_{n+1} \in B$ (cf. [8, §80C]). Let $f : B \rightarrow E$ be an algebraic map of degree n , and let $\overline{B}$ be the additive group generated by the elements of B . According to Dress [11, Proposition 1.1], there is a unique map $\overline{f} : \overline{B} \rightarrow E$ extending f , and $\overline{f}$ is also algebraic of degree n (see also [8, (80.44) Theorem (*Dress*)]). Assume further that $\overline{B}$ and E are commutative rings and B is closed under multiplication. If $f : B \rightarrow E$ is multiplicative, then the unique extension $\overline{f} : \overline{B} \rightarrow E$ of f to $\overline{B}$ is also multiplicative (cf. [8, (80.47) Theorem]).

Definition 3.16 Let $H \leq G$, and let $T_0, T_1, \dots, T_i$ be (H, A) -sets, where i is an integer with $0 \leq i \leq |G : H| + 1$. We define a (G, A) -set $\widehat{\text{Map}}_H(G, T_0, T_1, \dots, T_i)$ to be the set consisting of all elements $\widehat{f}$ of $\widehat{\text{Map}}_H(G, T_0 \dot{\cup} T_1 \dot{\cup} \dots \dot{\cup} T_i)$ containing $f \in \text{Map}_H(G, T_0 \dot{\cup} T_1 \dot{\cup} \dots \dot{\cup} T_i)$ such that $|\text{Im } f \cap T_\ell| \neq 0$ whenever $\ell \neq 0$ with the left action of G and the right action of A given by

$$g\widehat{f} = \widehat{gf} \quad \text{and} \quad \widehat{f}a = \widehat{f}_a$$

for all $g \in G$, $a \in A$, and $\widehat{f} \in \widehat{\text{Map}}_H(G, T_0, T_1, \dots, T_i)$.

Under the notation of Definition 3.16, we have

$$\widehat{\text{Map}}_H(G, T_0, T_1, \dots, T_i) = \begin{cases} \widehat{\text{Map}}_H(G, T_0) & \text{if } i = 0, \\ \widehat{\text{Map}}_H(G, T_1) & \text{if } T_0 = \emptyset \text{ and } i = 1, \\ \emptyset & \text{if } i = |G : H| + 1. \end{cases}$$

Proposition 3.17 For each $H \leq G$, the map $\widehat{\text{Map}}_H(G, -) : \Omega(H, A)^+ \rightarrow \Omega(G, A)$ is algebraic of degree $|G : H|$.

This proposition is analogous to [8, (80.43) Proposition (*Dress*)], and is an immediate consequence of the following lemma.

Lemma 3.18 Keep the notation of Definition 3.16, and assume further that $i \geq 1$. Set $\Theta_i = D_{[T_i]} \cdots D_{[T_1]} \widehat{\text{Map}}_H(G, -)$. Then

$$\Theta_i([T_0]) = [\widehat{\text{Map}}_H(G, T_0, T_1, \dots, T_i)].$$

Proof. The assertion is proved by an argument analogous to that in the proof of [8, (80.43) Proposition (*Dress*)]. $\square$

Tensor induction is multiplicative.

Lemma 3.19 For each $H \leq G$,

$$\widehat{\text{Map}}_H(G, T_1 \otimes T_2) \simeq \widehat{\text{Map}}_H(G, T_1) \otimes \widehat{\text{Map}}_H(G, T_2)$$

for all (H, A) -sets T_1 and T_2 .

Proof. If $f \in \text{Map}_H(G, T_1 \otimes T_2)$, then by Lemma 3.7, there exists a unique element $\widehat{\Psi_1(f)} \otimes \widehat{\Psi_2(f)}$ of $\widehat{\text{Map}_H(G, T_1)} \otimes \widehat{\text{Map}_H(G, T_2)}$, where $\Psi_i(f) \in \text{Map}_H(G, T_i)$ with $i = 1, 2$, such that

$$f(g) = \Psi_1(f)(g) \otimes \Psi_2(f)(g)$$

for all $g \in G$. Obviously, $\widehat{\Psi_1(f)} \otimes \widehat{\Psi_2(f)} = \widehat{\Psi_1(f')} \otimes \widehat{\Psi_2(f')}$ whenever $f \sim_A f'$. We now define a map $\widehat{\Psi} : \widehat{\text{Map}_H(G, T_1 \otimes T_2)} \rightarrow \widehat{\text{Map}_H(G, T_1)} \otimes \widehat{\text{Map}_H(G, T_2)}$ by

$$\widehat{f} \mapsto \widehat{\Psi_1(f)} \otimes \widehat{\Psi_2(f)}$$

for all $f \in \text{Map}_H(G, T_1 \otimes T_2)$. Observe that this map is (G, A) -equivariant and surjective. Moreover, by Remark 3.8,

$$|\widehat{\text{Map}_H(G, T_1 \otimes T_2)}| = (|T_1/A| \cdot |T_2/A|)^{|G/H|} \cdot |A| = |\widehat{\text{Map}_H(G, T_1)} \otimes \widehat{\text{Map}_H(G, T_2)}|.$$

Hence $\widehat{\Psi}$ is an isomorphism of (G, A) -sets. This completes the proof. $\square$

Combining Proposition 3.17 and Lemma 3.19 with [8, (80.47) Theorem], we obtain a result analogous to [8, (80.48) Theorem (*Dress*)].

Proposition 3.20 *For any $H \leq G$, there is a unique multiplicative map*

$$\overline{\text{Map}_H}(G, -) : \Omega(H, A) \rightarrow \Omega(G, A), \quad x \mapsto \overline{\text{Map}_H}(G, x)$$

extending $\widehat{\text{Map}_H}(G, -)$, called multiplicative induction or tensor induction, and this map is algebraic of degree $|G : H|$.

Remark 3.21 The multiplicative induction map $\overline{\text{Map}_H}(G, -) : \Omega(H, A) \rightarrow \Omega(G, A)$ with $A = \{\epsilon_A\}$ is introduced by Dress [11, §4].

Our concern is an explicit description of each element of $\text{Im } \overline{\text{Map}_H}(G, -)$ with $H \leq G$, and is to prove Eq.(1.1) (see also [8, (80.49) Corollary]).

Proposition 3.22 *Let $H \leq G$. For any (H, A) -sets T_0 and T ,*

$$\overline{\text{Map}_H}(G, [T_0] - [T]) = \sum_{i=0}^n (-1)^i [\widehat{\text{Map}_H}(G, T_0, T_1, \dots, T_i)],$$

where $n = |G : H|$ and $T = T_1 = \dots = T_n$.

Proof. We set $D_{[T]}^0 \Theta = \Theta = \overline{\text{Map}_H}(G, -) : \Omega(H, A) \rightarrow \Omega(G, A)$, and define inductively $D_{[T]}^i \Theta : \Omega(H, A) \rightarrow \Omega(G, A)$, $i = 1, 2, \dots$, by $D_{[T]}^i \Theta = D_{[T]}(D_{[T]}^{i-1} \Theta)$. From [8, (80.45)], we know that $\Theta([T_0] - [T]) = \sum_{i=0}^{\infty} (-1)^i D_{[T]}^i \Theta([T_0])$. Hence the assertion follows from Proposition 3.17 and Lemma 3.18. This completes the proof. $\square$

Remark 3.23 Let $(H, \sigma) \in \mathcal{S}(G, A)$. By Lemma 3.7 and Proposition 3.22, we can describe the structure of $\overline{\text{Map}}_H(G, -[(H/H)_\sigma])$. For each $X \in G\text{-set}$, let $\tilde{\Lambda}_{P(X)}$ be the reduced Lefschetz invariant of the poset $P(X)$ consisting of non-empty and proper subsets of X , which is an element of the Burnside ring $\Omega(G)$ (cf. [5, 29]). When $A = \{\epsilon_A\}$, $\overline{\text{Map}}_H(G, -[(H/H)_{1_H}])$ is identified with $\tilde{\Lambda}_{P(G/H)}$.

There is a Mackey decomposition formula which generalizes [32, §3(G.5)] (see also [2, Proposition 9.5]).

Proposition 3.24 *Let $H, K \leq G$. For each $x \in \Omega(H, A)$,*

$$\text{res}_K^G(\overline{\text{Map}}_H(G, x)) = \prod_{KgH \in K \backslash G/H} \overline{\text{Map}}_{K \cap gH}(K, \text{res}_{K \cap gH}^{gH} \circ \text{con}_H^g(x)).$$

Proof. By [8, (80.44) Theorem (*Dress*)], $\text{res}_K^G \circ \overline{\text{Map}}_H(G, -) : \Omega(H, A) \rightarrow \Omega(K, A)$ is the unique map extending the algebraic map

$$\begin{aligned} \text{res}_K^G \circ \widehat{\text{Map}}_H(G, -) &= \prod_{KgH \in K \backslash G/H} \widehat{\text{Map}}_{K \cap gH}(K, -) \circ \text{res}_{K \cap gH}^{gH} \circ \text{con}_H^g \\ &: \Omega(H, A)^+ \rightarrow \Omega(K, A), \end{aligned}$$

and so is $\prod_{KgH \in K \backslash G/H} \overline{\text{Map}}_{K \cap gH}(K, -) \circ \text{res}_{K \cap gH}^{gH} \circ \text{con}_H^g : \Omega(H, A) \rightarrow \Omega(K, A)$ (see [11, Proposition 1.2] and Propositions 3.9, 3.17, and 3.20). Thus the assertion holds. $\square$

4 The mark homomorphism

4A The first cohomology group

Following [12, §2], we provide preliminaries of the mark homomorphism for $\Omega(G, A)$ which is given in §4B.

Let $H \leq G$. The set $Z^1(H, A)$ is a right A -set with the action of A given in Definition 2.4, and is an abelian group with the product operation given by

$$\sigma \cdot \tau(h) = \sigma(h)\tau(h)$$

for all $\sigma, \tau \in Z^1(H, A)$ and $h \in H$. Obviously, the identity of $Z^1(H, A)$ is 1_H .

For each $\sigma \in Z^1(H, A)$, we denote by $\bar{\sigma}$ the A -orbit $\{\sigma^a \mid a \in A\}$ containing σ . Given $\sigma, \tau \in Z^1(H, A)$ and $a, b \in A$, it is easily seen that $\overline{\sigma^a \cdot \tau^b} = \overline{(\sigma \cdot \tau)^{ab}} = \bar{\sigma} \cdot \bar{\tau}$.

Definition 4.1 For each $H \leq G$, we define

$$H^1(H, A) := \{\bar{\sigma} \mid \sigma \in Z^1(H, A)\},$$

the set of A -orbits on $Z^1(H, A)$, and make it into an abelian group by defining

$$\bar{\sigma} \cdot \bar{\tau} = \overline{\sigma \cdot \tau}$$

for all $\sigma, \tau \in Z^1(H, A)$. (This product operation is well-defined.)

Let $H \leq G$. We denote by $\mathbb{Z}H^1(H, A)$ the group ring of $H^1(H, A)$ over $\mathbb{Z}$. Given $K \leq H$ and $g \in G$, there are ring homomorphisms

$$\begin{aligned} \text{con}_H^g : \mathbb{Z}H^1(H, A) &\rightarrow \mathbb{Z}H^1({}^gH, A), & \sum_{\bar{\sigma} \in H^1(H, A)} \ell_{\bar{\sigma}} \bar{\sigma} &\mapsto \sum_{\bar{\sigma} \in H^1(H, A)} \ell_{\bar{\sigma}} \overline{g\sigma} \quad \text{and} \\ \text{res}_K^H : \mathbb{Z}H^1(H, A) &\rightarrow \mathbb{Z}H^1(K, A), & \sum_{\bar{\sigma} \in H^1(H, A)} \ell_{\bar{\sigma}} \bar{\sigma} &\mapsto \sum_{\bar{\sigma} \in H^1(H, A)} \ell_{\bar{\sigma}} \overline{\sigma|_K}, \end{aligned}$$

where $\ell_{\bar{\sigma}} \in \mathbb{Z}$ with $\sigma \in Z^1(H, A)$ (see §2B), which are called the conjugation map and the restriction map, respectively (cf. [12, §2.2]). Obviously, the restriction map is well-defined. Let $\sigma \in Z^1(H, A)$. By Lemma 2.5, we have $\overline{g(\sigma^a)} = \overline{g\sigma}$ for any $a \in A$. Thus the conjugation map is well-defined.

Let Y be a (G, A) -set. The set of A -orbits yA , $y \in Y$, on Y is a left G -set. For each $y \in Y$, we denote by G_{yA} the stabilizer of the A -orbit yA in G , that is,

$$G_{yA} = \{g \in G \mid gy = ya \text{ for some } a \in A\},$$

and define a 1-cocycle $\sigma_y : G_{yA} \rightarrow A$ by

$$gy = y\sigma_y(g)$$

for all $g \in G_{yA}$. Obviously, $G_{(ya)A} = G_{yA}$ and $\sigma_{ya} = \sigma_y^a$ for any $y \in Y$ and $a \in A$.

Definition 4.2 Let Y be a (G, A) -set, and let $H \leq G$. We define

$$\text{inv}_H^A(Y) := \{y \in Y \mid H \leq G_{yA}\},$$

which is viewed as an (H, A) -subset of $\text{res}_H^G(Y)$, and define

$$[Y]_H := \frac{1}{|A|} \sum_{y \in \text{inv}_H^A(Y)} \text{res}_H^{G_{yA}}(\bar{\sigma}_y) = \sum_{yA \in \text{inv}_H^A(Y)/A} \text{res}_H^{G_{yA}}(\bar{\sigma}_y) \in \mathbb{Z}H^1(H, A).$$

Let Y_1 and Y_2 be (G, A) -sets, and let $H \leq G$. Obviously,

$$[Y_1 \dot{\cup} Y_2]_H = [Y_1]_H + [Y_2]_H.$$

Let $(y_1, y_2) \in Y_1 \times Y_2$. Given $g \in G$ and $a \in A$, $g(y_1 \otimes y_2) = (y_1 \otimes y_2)a$ if and only if $(gy_1b^{-1}, gy_2b) = (y_1, y_2a)$ for some $b \in A$. Hence we have

$$G_{(y_1 \otimes y_2)A} = G_{y_1A} \cap G_{y_2A} \quad \text{and} \quad \sigma_{y_1 \otimes y_2} = \sigma_{y_1}|_{G_{(y_1 \otimes y_2)A}} \cdot \sigma_{y_2}|_{G_{(y_1 \otimes y_2)A}}$$

(cf. [12, §2.3]). Moreover, $y_1 \otimes y_2 \in \text{inv}_H^A(Y_1 \otimes Y_2)$ if and only if $y_1 \in \text{inv}_H^A(Y_1)$ and $y_2 \in \text{inv}_H^A(Y_2)$. This means that

$$\begin{aligned} [Y_1]_H \cdot [Y_2]_H &= \frac{1}{|A|} \left(\sum_{y \in \text{inv}_H^A(Y_1)} \text{res}_H^{G_{yA}}(\overline{\sigma_y}) \right) \left(\sum_{yA \in \text{inv}_H^A(Y_2)/A} \text{res}_H^{G_{yA}}(\overline{\sigma_y}) \right) \\ &= \frac{1}{|A|} \sum_{y_1 \otimes y_2 \in \text{inv}_H^A(Y_1 \otimes Y_2)} \text{res}_H^{G_{(y_1 \otimes y_2)A}}(\overline{\sigma_{y_1 \otimes y_2}}) \\ &= [Y_1 \otimes Y_2]_H. \end{aligned}$$

Given $H \leq G$, we define a ring homomorphism $\rho_G^H : \Omega(G, A) \rightarrow \mathbb{Z}H^1(H, A)$ by

$$[Y] \mapsto [Y]_H$$

for all (G, A) -sets Y (cf. [12, §2.4]).

The ring homomorphisms $\rho_G^H : \Omega(G, A) \rightarrow \mathbb{Z}H^1(H, A)$ for $H \leq G$ form the map

$$\prod_{H \leq G} \rho_G^H : \Omega(G, A) \rightarrow \prod_{H \leq G} \mathbb{Z}H^1(H, A), \quad x \mapsto (\rho_G^H(x))_{H \leq G}$$

(cf. [12, §2.5]), which is injective (cf. [12, Theorem 1]).

4B The ghost ring

We continue reviewing part of [12, §2.4, §2.5], and define a ring monomorphism $\rho_G : \Omega(G, A) \rightarrow \mathcal{U}(G, A)$, $x \mapsto \prod_{H \leq G} \rho_G^H(x)$ (see Eq.(4.2)).

Definition 4.3 Let Y be a (G, A) -set, and let $(H, \sigma) \in \mathcal{S}(G, A)$. We define a subset $\text{inv}_{(H, \sigma)}(Y)$ of Y to be the set of $F_{(H, \sigma)}$ -invariants in Y , so that

$$\text{inv}_{(H, \sigma)}(Y) = \{y \in Y \mid hy = y\sigma(h) \text{ for all } h \in H\} = \{y \in \text{inv}_H^A(Y) \mid \sigma_y|_H = \sigma\},$$

and denote by A_σ the stabilizer $\{a \in A \mid \sigma = \sigma^a\}$ of $\sigma \in Z^1(H, A)$ in A .

Under the notation of Definition 4.3, the set $\text{inv}_{(H, \sigma)}(Y)$ is a free right A_σ -set with the action inherited from that of A on Y . For each $(H, \sigma) \in \mathcal{S}(G, A)$, we denote by $\text{inv}_{(H, \sigma)}(Y)/A_\sigma$ the set of A_σ -orbits on $\text{inv}_{(H, \sigma)}(Y)$.

Lemma 4.4 Let Y be a (G, A) -set, and let $H \leq G$. Then

$$[Y]_H = \sum_{\bar{\sigma} \in H^1(H, A)} |\text{inv}_{(H, \sigma)}(Y)/A_\sigma| \cdot \bar{\sigma}.$$

Moreover, $|\text{inv}_{(gH, g\sigma)}(Y)/A_{g\sigma}| = |\text{inv}_{(H, \sigma)}(Y)/A_\sigma|$ for any $\sigma \in Z^1(H, A)$ and $g \in G$.

Proof. The second statement is clear. To prove the first statement, we set

$$(Y/A)_{(H,\sigma)} = \{yA \in \text{inv}_H^A(Y)/A \mid \text{res}_H^{G_{yA}}(\overline{\sigma_y}) = \overline{\sigma}\}$$

for each $\sigma \in Z^1(H, A)$, so that

$$[Y]_H = \sum_{\overline{\sigma} \in H^1(H, A)} |(Y/A)_{(H,\sigma)}| \cdot \overline{\sigma}.$$

Hence it suffices to verify that $|(Y/A)_{(H,\sigma)}| = |\text{inv}_{(H,\sigma)}(Y)/A_\sigma|$ for any $\sigma \in Z^1(H, A)$. Let $\sigma \in Z^1(H, A)$. We make the set $\text{inv}_H^A(Y)$ into a free right A_σ -set by restriction of operators from A to A_σ . By definition,

$$\text{inv}_{(H,\sigma)}(Y)/A_\sigma = \{yA_\sigma \in \text{inv}_H^A(Y)/A_\sigma \mid \sigma_y|_H = \sigma\},$$

where $\text{inv}_H^A(Y)/A_\sigma$ is the set of A_σ -orbits $yA_\sigma := \{ya \mid a \in A_\sigma\}$, $y \in \text{inv}_H^A(Y)$, on $\text{inv}_H^A(Y)$. Let $y \in \text{inv}_H^A(Y)$, and suppose that $\sigma_y|_H = \sigma^a = \sigma^b$ for some $a, b \in A$. Then $ab^{-1} \in A_\sigma$ and $ya^{-1}A_\sigma = yb^{-1}A_\sigma \in \text{inv}_{(H,\sigma)}(Y)/A_\sigma$. (Note that $\sigma_{(yc)}|_H = \sigma^{ac}$ for any $c \in A$.) Hence there is a bijection $(Y/A)_{(H,\sigma)} \rightarrow \text{inv}_{(H,\sigma)}(Y)/A_\sigma$ given by

$$yA \mapsto ya^{-1}A_\sigma,$$

where $\sigma_y|_H = \sigma^a$ with $a \in A$, for all $yA \in (Y/A)_{(H,\sigma)}$. This completes the proof. $\square$

The following lemma is [27, Lemma 3.3].

Lemma 4.5 *Let $(H, \sigma), (U, \tau) \in \mathcal{S}(G, A)$. Then*

$$|\text{inv}_{(H,\sigma)}((G/U)_\tau)/A_\sigma| = |\{gU \in G/U \mid H \leq {}^gU \text{ and } (g\tau)|_H = {}_A\sigma\}|.$$

Let $H, U \leq G$, and consider G/U to be a left G -set with the action of G given by the product operation on G . Following [8, (80.5) Proposition], we define

$$\text{inv}_H(G/U) := \{gU \in G/U \mid H \leq {}^gU\}. \quad (4.1)$$

Lemma 4.6 (a) *Let $H \leq G$, and let $(U, \tau) \in \mathcal{S}(G, A)$. Then*

$$[(G/U)_\tau]_H = \sum_{gU \in \text{inv}_H(G/U)} \text{res}_H^{gU} \circ \text{con}_U^g(\overline{\tau}).$$

(b) *Let $K \leq H \leq G$, and let $(U, \tau) \in \mathcal{S}(H, A)$. Then for any $r \in G$,*

$$[{}^r((H/U)_\tau)]_{rK} = \text{con}_K^r([(H/U)_\tau]_K).$$

If $H = G$, then for any $r \in G$,

$$[(G/U)_\tau]_{rK} = \text{con}_K^r([(G/U)_\tau]_K).$$

Proof. (a) Although the assertion follows from Lemmas 4.4 and 4.5, we directly prove it. In the proof of Lemma 4.4, if $Y = (G/U)_\tau$, then by Lemmas 2.5 and 2.6,

$$(Y/A)_{(H,\sigma)} = \{(\epsilon_A, gU)A \in \text{inv}_H^A(Y)/A \mid \text{res}_H^{gU} \circ \text{con}_U^g(\bar{\tau}) = \bar{\sigma}\},$$

whence

$$[(G/U)_\tau]_H = \sum_{\bar{\sigma} \in H^1(H,A)} |(Y/A)_{(H,\sigma)}| \cdot \bar{\sigma} = \sum_{gU \in \text{inv}_H(G/U)} \text{res}_H^{gU} \circ \text{con}_U^g(\bar{\tau}).$$

(b) By Lemma 2.10, it suffices to prove the first assertion. We have

$$\begin{aligned} [({}^rH/{}^rU)_{r\tau}]_{rK} &= \sum_{h'{}^rU \in \text{inv}_{rK}({}^rH/{}^rU)} \text{res}_{rK}^{h'{}^rU} \circ \text{con}_{rU}^{h'}(\overline{r\tau}) \\ &= \sum_{h'{}^rU \in \text{inv}_{rK}({}^rH/{}^rU)} \text{con}_{rK}^r \circ \text{res}_K^{r^{-1}h'{}^rU} \circ \text{con}_U^{r^{-1}h'}(\bar{\tau}) \\ &= \text{con}_K^r([(H/U)_\tau]_K). \end{aligned}$$

Hence the first assertion follows from Lemma 2.10. This completes the proof. $\square$

Definition 4.7 We define

$$\mathfrak{U}(G, A) := \left\{ (x_H)_{H \leq G} \in \prod_{H \leq G} \mathbb{Z}H^1(H, A) \mid \text{con}_H^g(x_H) = x_{gH} \text{ for all } g \in G \right\},$$

the ghost ring of $\Omega(G, A)$, which is a subring of $\prod_{H \leq G} \mathbb{Z}H^1(H, A)$.

Remark 4.8 The family of $\mathbb{Z}$ -algebras $\mathbb{Z}H^1(H, A)$ for $H \leq G$, together with conjugation maps and restriction maps, defines a $\mathbb{Z}$ -algebra restriction functor $\mathbb{Z}H^1(-, A)$ defined in [4, 1.1. Definition]. The rings $\Omega(G, A)$ and $\mathfrak{U}(G, A)$ are identified with $\mathbb{Z}H^1(G, A)_+$ and $\mathbb{Z}H^1(G, A)^+$, respectively, which are obtained by the plus constructions $\mathbb{Z}H^1(-, A) \mapsto \mathbb{Z}H^1(-, A)_+$ and $\mathbb{Z}H^1(-, A) \mapsto \mathbb{Z}H^1(-, A)^+$; moreover, the Green functor given in Proposition 2.16 is identified with $\mathbb{Z}H^1(-, A)_+$ (see [4]).

From Proposition 2.14 and Lemma 4.6, we know that there is an additive map $\rho_G : \Omega(G, A) \rightarrow \mathfrak{U}(G, A)$ given by

$$[(G/U)_\tau] \mapsto \left(\sum_{gU \in \text{inv}_H(G/U)} \text{res}_H^{gU} \circ \text{con}_U^g(\bar{\tau}) \right)_{H \leq G}$$

for all $(U, \tau) \in \mathcal{R}(G, A)$ (cf. [4, 2.3.]), which is called the mark homomorphism. Since

$$\rho_G([Y]) = ([Y]_H)_{H \leq G}$$

for all (G, A) -sets Y , the mark homomorphism is a ring homomorphism defined by

$$\rho_G(x) = (\rho_G^H(x))_{H \leq G} \quad (4.2)$$

for all $x \in \Omega(G, A)$ (cf. [12, §2.5]). We write $\rho = \rho_G$ for shortness' sake.

According to [4, (2.3a)], there is a map $\eta : \mathfrak{U}(G, A) \rightarrow \Omega(G, A)$ given by

$$\left(\sum_{\bar{\sigma} \in H^1(H, A)} \ell_{(H, \bar{\sigma})} \bar{\sigma} \right)_{H \leq G} \mapsto \sum_{H \leq G} \sum_{U \leq H} |U| \mu(U, H) \sum_{\bar{\sigma} \in H^1(H, A)} \ell_{(H, \bar{\sigma})} [(G/U)_{\sigma|_U}]$$

for all $\ell_{(H, \bar{\sigma})} \in \mathbb{Z}$ with $H \leq G$ and $\sigma \in Z^1(H, A)$.

We quote concise versions of [4, 2.4. Proposition] and [12, Theorem 1].

Proposition 4.9 (a) $\eta \circ \rho = |G| \text{id}_{\Omega(G, A)}$. (b) $\rho \circ \eta = |G| \text{id}_{\mathfrak{U}(G, A)}$.

Corollary 4.10 *The mark homomorphism ρ is injective.*

4C Invariant of tensor induction

Let $H \leq G$. By Example 3.13 and Proposition 3.14, we have

$$\rho([\widehat{\text{Map}}_H(G, (H/H)_\sigma)]) = (\overline{\sigma^{\otimes G}}|_K)_{K \leq G} = \left(\prod_{KgH \in K \backslash G/H} \overline{(g\sigma)|_{K \cap gH}^{\otimes K}} \right)_{K \leq G} \quad (4.3)$$

for all $\sigma \in Z^1(H, A)$. Let T be an (H, A) -set. We are interested in the description of $\rho([\widehat{\text{Map}}_H(G, T)])$, which naturally extends Eq.(4.3) (see Proposition 4.14). For each $K \leq G$, the K -component $[\widehat{\text{Map}}_H(G, T)]_K$ of $\rho([\widehat{\text{Map}}_H(G, T)])$ is also associated with a Mackey decomposition formula (see Proposition 3.9).

Let Y be a (G, A) -set, and let $K \leq G$. By Definition 4.2,

$$[Y]_K = \frac{1}{|A|} \sum_{y \in \text{inv}_K^A(Y)} \text{res}_K^{G_{yA}}(\overline{\sigma_y}) = \sum_{yA \in \text{inv}_K^A(Y)/A} \text{res}_K^{G_{yA}}(\overline{\sigma_y}).$$

Concerning this formula, we have

$$[Y]_K = \frac{1}{|A|} \sum_{y \in \text{inv}_K^A(Y)} \text{res}_K^{G_{yA}}(\overline{\sigma_y}) = \frac{1}{|A|} \sum_{y \in \text{inv}_K^A(\text{res}_K^G(Y))} \overline{\sigma_y} = [\text{res}_K^G(Y)]_K. \quad (4.4)$$

Obviously, this fact implies that $\rho_G^K(x) = \rho_K^K(\text{res}_K^G(x))$ for any $x \in \Omega(G, A)$ which is applied to the following lemma.

Lemma 4.11 *Let $H, K \leq G$. For any $x \in \Omega(H, A)$,*

$$\rho_G^K(\overline{\text{Map}}_H(G, x)) = \prod_{KgH \in K \backslash G/H} \rho_K^K(\overline{\text{Map}}_{K \cap {}^g H}(K, \text{res}_{K \cap {}^g H}^{gH} \circ \text{con}_H^g(x))).$$

Proof. Since $\rho_G^K(\overline{\text{Map}}_H(G, x)) = \rho_K^K(\text{res}_K^G(\overline{\text{Map}}_H(G, x)))$ for any $x \in \Omega(H, A)$, the assertion follows from Proposition 3.24. This completes the proof. $\square$

Definition 4.12 Let $H \leq G$. We define a map $-\otimes^G : \mathbb{Z}H^1(H, A) \rightarrow \mathbb{Z}H^1(G, A)$ by

$$\sum_{\bar{\sigma} \in H^1(H, A)} \ell_{\bar{\sigma}} \bar{\sigma} \mapsto \left(\sum_{\bar{\sigma} \in H^1(H, A)} \ell_{\bar{\sigma}} \bar{\sigma} \right)^{\otimes G} := \sum_{\bar{\sigma} \in H^1(H, A)} \ell_{\bar{\sigma}} \overline{\sigma}^{\otimes G}$$

for all $\ell_{\bar{\sigma}} \in \mathbb{Z}$ with $\sigma \in Z^1(H, A)$. (This map is well-defined; see Remark 3.12.)

Lemma 4.13 *Let $H \leq G$, and let T be an (H, A) -set. Then*

$$[\widehat{\text{Map}}_H(G, T)]_G = \frac{1}{|A|} \sum_{\widehat{f} \in \text{inv}_G^A(\widehat{\text{Map}}_H(G, T))} \overline{\sigma}_{\widehat{f}} = \frac{1}{|A|} \sum_{t \in \text{inv}_H^A(T)} \overline{\sigma}_t^{\otimes G} = [T]_H^{\otimes G}.$$

Proof. Fix a complete set $\{g_1, g_2, \dots, g_n\}$ with $g_1 = \epsilon$ of representatives of G/H . Let $t \in \text{inv}_H^A(T)$. We define an element $f_{(t)}$ of $\text{Map}_H(G, T)$ by

$$f_{(t)}(g_j^{-1}) = t$$

for all $j \in [n]$. For any $g \in G$ and $j \in [n]$, if $g_j H = gg_{j'} H$, then

$$(gf_{(t)})(g_j^{-1}) = (g_j^{-1}gg_{j'})f_{(t)}(g_j^{-1}) = (g_j^{-1}gg_{j'})t = f_{(t)}(g_j^{-1})\sigma_t(g_j^{-1}gg_{j'}).$$

This, combined with Lemma 3.7, shows that $\widehat{gf_{(t)}} = \widehat{f_{(t)}}(\sigma_t^{\otimes G})(g)$ for all $g \in G$ (see Definition 3.11). Hence $G = G_{\widehat{f_{(t)}}A}$, $\widehat{f_{(t)}} \in \text{inv}_G^A(\widehat{\text{Map}}_H(G, T))$, and $\sigma_{\widehat{f_{(t)}}} = \sigma_t^{\otimes G}$.

We now define a map $\widehat{\Gamma} : \text{inv}_H^A(T)/A \rightarrow \text{inv}_G^A(\widehat{\text{Map}}_H(G, T))/A$ by

$$\widehat{\Gamma}(tA) = \widehat{f_{(t)}}A$$

for all $t \in \text{inv}_H^A(T)$. This map is well-defined, because, by Lemma 3.7, $\widehat{f_{(ta)}} = \widehat{f_{(t)}}b$ with $b = {}^{g_1}a {}^{g_2}a \dots {}^{g_n}a$ for any $a \in A$. If $\widehat{\Gamma}(t_1 A) = \widehat{\Gamma}(t_2 A)$ with $t_1, t_2 \in \text{inv}_H^A(T)$, then $\widehat{f_{(t_1)}} = \widehat{f_{(t_2)}}a$ for some $a \in A$, and hence $t_1 = t_2 b$ for some $b \in A$. Thus $\widehat{\Gamma}$ is injective. Let $\widehat{f} \in \text{inv}_G^A(\widehat{\text{Map}}_H(G, T))$ with $f \in \text{Map}_H(G, T)$, and let $g \in G$. Given $j \in [n]$, we have $(gf)(g_j^{-1}) = f(g_j^{-1})a_j(g)$ for some $a_j(g) \in A$. Set $t = f(\epsilon)$. Then

$$ht = hf(\epsilon) = f(h) = (hf)(\epsilon) = ta_1(h)$$

for all $h \in H$, which yields $t \in \text{inv}_H^A(T)$. Observe now that for any $j \in [n]$,

$$f(g_j^{-1})a_j(g_j) = (g_j f)(g_j^{-1}) = f(\epsilon) = t = f_{(t)}(g_j^{-1}).$$

By Lemma 3.7, we have $\widehat{f} = \widehat{f_{(t)}}a$, where $a = ({}^g a_1(g_1) {}^{g^2} a_2(g_2) \cdots {}^{g^n} a_n(g_n))^{-1}$, so that $\widehat{\Gamma}(tA) = \widehat{f}A$. Thus $\widehat{\Gamma}$ is bijective. The assertion now follows from the fact that $\sigma_{\widehat{f_{(t)}}} = \sigma_t^{\otimes G}$ for all $t \in \text{inv}_H^A(T)$. This completes the proof. $\square$

The following proposition generalizes the equation in [32, p. 39] (see also [2, Lemma 9.2], [9, p. 149], and [30, p. 111, Eq.(2)]).

Proposition 4.14 *Let $H, K \leq G$. For each (H, A) -set T ,*

$$[\widehat{\text{Map}}_H(G, T)]_K = \prod_{KgH \in K \backslash G/H} [{}^g T]_{K \cap {}^g H}^{\otimes K}.$$

Proof. Combining Lemma 4.13 with Lemma 4.11, we have

$$\begin{aligned} [\widehat{\text{Map}}_H(G, T)]_K &= \prod_{KgH \in K \backslash G/H} [\widehat{\text{Map}}_{K \cap {}^g H}(K, \text{res}_{K \cap {}^g H}^{{}^g H}({}^g T))]_K \\ &= \prod_{KgH \in K \backslash G/H} [\text{res}_{K \cap {}^g H}^{{}^g H}({}^g T)]_{K \cap {}^g H}^{\otimes K}. \end{aligned}$$

Hence the assertion follows from Eq.(4.4). This completes the proof. $\square$

How about the description of $\rho(\overline{\text{Map}}_H(G, x))$ for any $H \leq G$ and $x \in \Omega(H, A)$? By using Eq.(1.1), we are successful in proving Eq.(1.2) (see Theorem 4.16).

Lemma 4.15 *Let $H \leq G$. For any (H, A) -sets T_0 and T ,*

$$\rho_G^G(\overline{\text{Map}}_H(G, [T_0] - [T])) = [\widehat{\text{Map}}_H(G, T_0)]_G - [\widehat{\text{Map}}_H(G, T)]_G.$$

Proof. We may assume that $H < G$. By Proposition 3.22,

$$\overline{\text{Map}}_H(G, [T_0] - [T]) = [\widehat{\text{Map}}_H(G, T_0)] + \sum_{i=1}^n (-1)^i [\widehat{\text{Map}}_H(G, T_0, T_1, \dots, T_i)],$$

where $n = |G : H|$ and $T = T_1 = \cdots = T_n$. If $i \in [n]$ and $i \geq 2$, then obviously, $[\widehat{\text{Map}}_H(G, T_0, T_1, \dots, T_i)]_G = 0$. Moreover, we have

$$\text{inv}_G^A(\widehat{\text{Map}}_H(G, T_0, T_1)) = \text{inv}_G^A(\widehat{\text{Map}}_H(G, \emptyset, T_1)) = \text{inv}_G^A(\widehat{\text{Map}}_H(G, T_1)),$$

completing the proof. $\square$

The following theorem, which is equivalent to Eq.(1.2), is an extension of Proposition 4.14 and is a generalization of [32, §3(b.3)].

Theorem 4.16 *Let $H \leq G$, and define a map $\text{jnd}_H^G : \mathcal{U}(H, A) \rightarrow \mathcal{U}(G, A)$ by*

$$(x_L)_{L \leq H} \mapsto \left(\prod_{KgH \in K \backslash G/H} \text{con}_{KgH}^g(x_{KgH})^{\otimes K} \right)_{K \leq G}$$

for all $(x_L)_{L \leq H} \in \mathcal{U}(H, A)$. Then the diagram

$$\begin{array}{ccc} \Omega(G, A) & \xrightarrow{\rho} & \mathcal{U}(G, A) \\ \overline{\text{Map}}_H(G, -) \uparrow & & \uparrow \text{jnd}_H^G \\ \Omega(H, A) & \xrightarrow{\rho_H} & \mathcal{U}(H, A) \end{array}$$

is commutative, where $\rho_H : \Omega(H, A) \rightarrow \mathcal{U}(H, A)$ is the mark homomorphism.

Proof. We prove Eq.(1.2). Let $x \in \Omega(H, A)$. We may assume that $x = [T_0] - [T]$ for some (H, A) -sets T_0 and T . Let $K \leq G$. Then by Lemmas 4.11 and 4.15, we have

$$\begin{aligned} & \rho_G^K(\overline{\text{Map}}_H(G, [T_0] - [T])) \\ &= \prod_{KgH \in K \backslash G/H} \rho_K^K(\overline{\text{Map}}_{K \cap gH}(K, [\text{res}_{K \cap gH}^{gH}({}^gT_0)] - [\text{res}_{K \cap gH}^{gH}({}^gT)])) \\ &= \prod_{KgH \in K \backslash G/H} \left\{ [\widehat{\text{Map}}_{K \cap gH}(K, \text{res}_{K \cap gH}^{gH}({}^gT_0))]_K \right. \\ & \quad \left. - [\widehat{\text{Map}}_{K \cap gH}(K, \text{res}_{K \cap gH}^{gH}({}^gT))]_K \right\}. \end{aligned}$$

Moreover, it follows from Eq.(4.4) and Lemma 4.13 that

$$\rho_G^K(\overline{\text{Map}}_H(G, [T_0] - [T])) = \prod_{KgH \in K \backslash G/H} \left\{ [{}^gT_0]_{K \cap gH}^{\otimes K} - [{}^gT]_{K \cap gH}^{\otimes K} \right\}. \quad (4.5)$$

By Lemma 4.6(b), $[{}^gT_1]_{K \cap gH} = \text{con}_{KgH}^g([T_1]_{KgH})$, where $T_1 = T_0$ or $T_1 = T$, for all $g \in G$. Hence Eq.(1.2) follows from Eq.(4.2). This completes the proof. $\square$

Remark 4.17 Given $(H, \sigma) \in \mathcal{S}(G, A)$, it follows from Lemma 4.6 and Eq.(4.5) that

$$\rho(\overline{\text{Map}}_H(G, -[(H/H)_\sigma])) = \left((-1)^{|K \backslash G/H|} \prod_{KgH \in K \backslash G/H} \overline{(g\sigma)_{K \cap gH}^{\otimes K}} \right)_{K \leq G}$$

(see also Eq.(4.3)). Here we return to Remark 3.23. Deducing this fact directly from Lemma 3.7 and Proposition 3.22 requires the use of [25, (24d)] which provides a combinatorial explanation. Let $H, K \leq G$. When $A = \{\epsilon_A\}$, the K -component of $\rho(\tilde{\Lambda}_{P(G/H)})$ is $(-1)^{|K \backslash G/H|}$ (see [29, Proposition 5.1] and [32, Lemma 3.6]).

For each $H \leq G$, we denote by $\Omega(H, A)^\times$ the unit group of $\Omega(H, A)$, and consider this abelian group as a $\mathbb{Z}$ -module. Note that the $\mathbb{Z}$ -module structure of $\Omega(H, A)^\times$ is different from that of $\Omega(H, A)$.

There is a fact relative to [2, Theorem 9.6] and [32, Lemma 3.1].

Theorem 4.18 *The family of $\mathbb{Z}$ -modules $\Omega(H, A)^\times$ for $H \leq G$, together with conjugation, restriction, and multiplicative induction maps inherited from those on the family of $\mathbb{Z}$ -algebras $\Omega(H, A)$ for $H \leq G$ defines a Mackey functor on G .*

Proof. Let $\text{jnd}_K^H : \Omega(K, A)^\times \rightarrow \Omega(H, A)^\times$ with $K \leq H \leq G$ be the map inherited from $\text{Map}_K(H, -) : \Omega(K, A) \rightarrow \Omega(H, A)$. By [4, 1.1. Definition], Lemma 2.10, and Proposition 3.24, it suffices to verify that for any $U \leq V \leq H \leq G$ and $g \in G$,

$$\text{con}_H^g \circ \text{jnd}_U^H = \text{jnd}_{gU}^{gH} \circ \text{con}_U^g \quad \text{and} \quad \text{jnd}_V^H \circ \text{jnd}_U^V = \text{jnd}_U^H. \quad (4.6)$$

Given $H \leq G$ and $g \in G$, we define a map $\text{con}_H^g : \mathcal{U}(H, A) \rightarrow \mathcal{U}(gH, A)$ by

$$(x_K)_{K \leq H} \mapsto (\text{con}_K^g(x_K))_{gK \leq gH}$$

for all $(x_K)_{K \leq H} \in \mathcal{U}(H, A)$. Let $U \leq V \leq H \leq G$, and let $g \in G$. Given $K \leq H$ and $(x_L)_{L \leq U} \in \mathcal{U}(U, A)$, we have

$$\text{con}_K^g(\text{con}_{K^h \cap U}^h(x_{K^h \cap U})^{\otimes K}) = (\text{con}_{(gK)^{g_h \cap gU}}^{g_h} \circ \text{con}_{K^h \cap U}^g(x_{K^h \cap U}))^{\otimes gK}$$

for all $h \in H$ and

$$\begin{aligned} \prod_{KhV \in K \setminus H/V} \text{con}_{K_h}^h \left(\prod_{K_h rU \in K_h \setminus V/U} \text{con}_{K_h rU}^r(x_{K_h rU})^{\otimes K_h} \right)^{\otimes K} \\ = \prod_{KhU \in K \setminus H/U} \text{con}_{K^h \cap U}^h(x_{K^h \cap U})^{\otimes K}, \end{aligned}$$

where $K_h = K^h \cap V$ (see Lemma 3.15). Relative to ‘jnd’ defined in Theorem 4.16, these equations enable us to obtain the equations

$$\text{con}_H^g \circ \text{jnd}_U^H = \text{jnd}_{gU}^{gH} \circ \text{con}_U^g \quad \text{and} \quad \text{jnd}_V^H \circ \text{jnd}_U^V = \text{jnd}_U^H.$$

By Lemma 4.6(b), $\text{con}_H^g \circ \rho_H = \rho_{gH} \circ \text{con}_H^g$ and $\text{con}_U^g \circ \rho_U = \rho_{gU} \circ \text{con}_U^g$. Hence Eq.(4.6) follows from Corollary 4.10 and Theorem 4.16. This completes the proof. $\square$

5 Fundamentals of monomial Burnside rings

5A The Burnside homomorphism

The discussion in this section is a special case of [28, §9] (see also [27, §3, §4]).

For each $(U, \tau) \in \mathcal{S}(G, A)$, we set

$$N_G(U, \tau) = \{g \in G \mid {}^g U = U \text{ and } \text{con}_U^g(\bar{\tau}) = \bar{\tau}\}.$$

By definition, the elements $(x_H^{(U, \tau)})_{H \leq G}$ for $(U, \tau) \in \mathcal{R}(G, A)$, where

$$x_H^{(U, \tau)} = \begin{cases} \sum_{gN_G(U, \tau) \in N_G(U)/N_G(U, \tau)} \text{con}_U^{rg}(\bar{\tau}) & \text{if } H = {}^r U \text{ with } r \in G, \\ 0 & \text{otherwise,} \end{cases}$$

form a free $\mathbb{Z}$ -basis of the ghost ring $\mathfrak{U}(G, A)$. We define

$$\tilde{\Omega}(G, A) := \coprod_{(K, \nu) \in \mathcal{R}(G, A)} \mathbb{Z},$$

so that there exists an isomorphism $\kappa : \tilde{\Omega}(G, A) \xrightarrow{\sim} \mathfrak{U}(G, A)$ of $\mathbb{Z}$ -lattices given by

$$(\delta_{(U, \tau)}(K, \nu))_{(K, \nu) \in \mathcal{R}(G, A)} \mapsto (x_H^{(U, \tau)})_{H \leq G}$$

for all $(U, \tau) \in \mathcal{R}(G, A)$, where δ is the Kronecker delta.

Definition 5.1 We define an additive map $\varphi : \Omega(G, A) \rightarrow \tilde{\Omega}(G, A)$ by

$$\varphi([(G/U)_\tau]) = (\text{inv}_{(K, \nu)}((G/U)_\tau)/A_\nu)_{(K, \nu) \in \mathcal{R}(G, A)}$$

for all $(U, \tau) \in \mathcal{R}(G, A)$ (see Lemma 4.5), and call it the Burnside homomorphism.

Proposition 5.2 *The diagram*

$$\begin{array}{ccc} \Omega(G, A) & \xrightarrow{\varphi} & \tilde{\Omega}(G, A) \\ & \searrow \rho & \downarrow \kappa \\ & & \mathfrak{U}(G, A) \end{array}$$

is commutative. In particular, the Burnside homomorphism φ is injective.

Proof. The assertion follows from Lemma 4.4 and Corollary 4.10. $\square$

Let $(U, \tau) \in \mathcal{R}(G, A)$. By Lemma 2.6, $N_G(U, \tau)$ contains U . Observe that for any $(K, \nu) \in \mathcal{R}(G, A)$, the (K, ν) -component of $\varphi([(G/U)_\tau])$ is divisible by $|N_G(U, \tau)/U|$ (see Lemma 4.5). We define

$$y^{(U, \tau)} := \frac{1}{|N_G(U, \tau)/U|} \varphi([(G/U)_\tau]) = \left(\frac{|\text{inv}_{(K, \nu)}((G/U)_\tau)/A_\nu|}{|N_G(U, \tau)/U|} \right)_{(K, \nu) \in \mathcal{R}(G, A)}.$$

Proposition 5.3 *The elements $y^{(U, \tau)}$ for $(U, \tau) \in \mathcal{R}(G, A)$ form a free $\mathbb{Z}$ -basis of the $\mathbb{Z}$ -lattice $\tilde{\Omega}(G, A)$.*

Proof. The proof is completely analogous to that of [8, (80.15) Proposition]. $\square$

5B The Cauchy-Frobenius homomorphism

We aim to state a fundamental theorem for the monomial Burnside ring $\Omega(G, A)$ (see Theorem 5.9).

Definition 5.4 For each $(U, \tau) \in \mathcal{S}(G, A)$, let $W_G(U, \tau)$ denote the factor group $N_G(U, \tau)/U$. We define

$$\text{Obs}(G, A) := \coprod_{(U, \tau) \in \mathcal{R}(G, A)} \mathbb{Z}/|W_G(U, \tau)|\mathbb{Z},$$

the obstruction group of $\Omega(G, A)$.

The following fact is a corollary to Proposition 5.3.

Corollary 5.5 $\tilde{\Omega}(G, A)/\text{Im}\varphi \simeq \text{Obs}(G, A)$.

Proof. The proof is completely analogous to that of [27, Corollary 3.8]. $\square$

Let p be a prime, and let $\mathbb{Z}_{(p)}$ be the localization of $\mathbb{Z}$ at p . For each $\mathbb{Z}$ -module M , we set $M_{(p)} = \mathbb{Z}_{(p)} \otimes_{\mathbb{Z}} M$ and $M_{(\infty)} = M$. Let $(U, \tau) \in \mathcal{S}(G, A)$. We denote by $W_G(U, \tau)_p$ a Sylow p -subgroup of $W_G(U, \tau)$, and set $W_G(U, \tau)_{\infty} = W_G(U, \tau)$.

Let p be a prime or the symbol ∞ hereafter. By Proposition 2.14, the elements $[(G/H)_{\sigma}]$ for $(H, \sigma) \in \mathcal{R}(G, A)$ form a free $\mathbb{Z}_{(p)}$ -basis of the $\mathbb{Z}_{(p)}$ -lattice $\Omega(G, A)_{(p)}$. We identify $\tilde{\Omega}(G, A)_{(p)}$ and $\text{Obs}(G, A)_{(p)}$ with

$$\coprod_{(K, \nu) \in \mathcal{R}(G, A)} \mathbb{Z}_{(p)} \quad \text{and} \quad \coprod_{(U, \tau) \in \mathcal{R}(G, A)} \mathbb{Z}_{(p)}/|W_G(U, \tau)_p|\mathbb{Z}_{(p)},$$

respectively. Let $\varphi^{(p)}$ denote the monomorphism from $\Omega(G, A)_{(p)}$ to $\tilde{\Omega}(G, A)_{(p)}$ determined by φ . (So $\varphi^{(\infty)} = \varphi$.) Then by Corollary 5.5, we have

$$\tilde{\Omega}(G, A)_{(p)}/\text{Im}\varphi^{(p)} \simeq \text{Obs}(G, A)_{(p)}. \quad (5.1)$$

The expression ‘ $x \bmod \ell$ ’ with $x, \ell \in \mathbb{Z}_{(p)}$ denotes the coset $x + \ell\mathbb{Z}_{(p)}$ of $\ell\mathbb{Z}_{(p)}$ in $\mathbb{Z}_{(p)}$ containing x . Let $(U, \tau) \in \mathcal{S}(G, A)$. Given $(y_{(H, \sigma)})_{(H, \sigma) \in \mathcal{R}(G, A)} \in \tilde{\Omega}(G, A)_{(p)}$, $y_{(U, \tau)}$ denotes $y_{(H, \sigma)}$ for a representative $(H, \sigma) \in \mathcal{R}(G, A)$ of the F -orbit on $\mathcal{S}(G, A)$ containing (U, τ) . For each $g \in N_G(U, \tau)$, we set

$$H_{\tau}^1(\langle g \rangle U, A) = \{\bar{\nu} \in H^1(\langle g \rangle U, A) \mid \text{res}_U^{\langle g \rangle U}(\bar{\nu}) = \bar{\tau}\}.$$

Definition 5.6 We define an additive map $\psi^{(p)} : \tilde{\Omega}(G, A)_{(p)} \rightarrow \text{Obs}(G, A)_{(p)}$ by

$$(y_{(K, \nu)})_{(K, \nu) \in \mathcal{R}(G, A)} \mapsto \left(\sum_{\substack{gU \in W_G(U, \tau)_p, \\ \bar{\nu} \in H_{\tau}^1(\langle g \rangle U, A)}} y_{(\langle g \rangle U, \nu)} \bmod |W_G(U, \tau)_p| \right)_{(U, \tau) \in \mathcal{R}(G, A)}$$

for all $(y_{(K,\nu)})_{(K,\nu) \in \mathcal{R}(G,A)} \in \tilde{\Omega}(G, A)_{(p)}$, and call it the Cauchy-Frobenius homomorphism.

Remark 5.7 (1) When p is a prime, $\psi^{(p)}$ is independent of the choice of a Sylow p -subgroup $W_G(U, \tau)_p$ of $W_G(U, \tau)$ (cf. [28, §9]). (2) When $p = \infty$, we write $\psi = \psi^{(\infty)}$.

For each $(H, \sigma) \in \mathcal{R}(G, A)$, it follows from Lemma 4.5 that

$$\psi^{(p)} \circ \varphi^{(p)}([(G/H)_\sigma]) = \left(\sum_{gU \in W_G(U, \tau)_p} |I_{gU, \tau}^{(H, \sigma)}| \bmod |W_G(U, \tau)_p| \right)_{(U, \tau) \in \mathcal{R}(G, A)}, \quad (5.2)$$

where

$$I_{gU, \tau}^{(H, \sigma)} = \{rH \in G/H \mid \langle g \rangle U \leq {}^r H \text{ and } (r\sigma)|_U = {}_A \tau\}.$$

The following lemma, which is a special case of [28, Lemma 9.2], is a consequence of the Cauchy-Frobenius lemma (see, e.g., [33, 2.7 Lemma]).

Lemma 5.8 *Let $(H, \sigma), (U, \tau) \in \mathcal{R}(G, A)$. For any $V \leq N_G(U, \tau)$ with $U \leq V$,*

$$\sum_{gU \in V/U} |I_{gU, \tau}^{(H, \sigma)}| \equiv 0 \pmod{|V/U|}.$$

Proof. The proof is analogous to that of [28, Lemma 9.2], and is also analogous to part of the proof of [27, Lemma 4.1]. $\square$

We are now in a position to show a special case of [28, Theorem 9.4], which is a generalization of [9, Proposition 1.3.5] and [32, Lemma 2.1].

Theorem 5.9 (Fundamental theorem) *The sequence*

$$0 \longrightarrow \Omega(G, A)_{(p)} \xrightarrow{\varphi^{(p)}} \tilde{\Omega}(G, A)_{(p)} \xrightarrow{\psi^{(p)}} \text{Obs}(G, A)_{(p)} \longrightarrow 0$$

of additive groups is exact.

Proof. By Proposition 5.2, $\varphi^{(p)}$ is injective. Moreover, it is easily verified that $\psi^{(p)}$ is surjective (see, e.g., the proof of [27, Lemma 4.3]). Using Eqs.(5.1) and (5.2) and Lemma 5.8, we have $\text{Im } \varphi^{(p)} = \text{Ker } \psi^{(p)}$, completing the proof. $\square$

5C Idempotents of Burnside rings

The Burnside ring $\Omega(G)$ of G , which is defined to be the Grothendieck ring of G -set, is the commutative unital ring consisting of all formal $\mathbb{Z}$ -linear combinations of the symbols $[G/H]$ for $H \in \mathcal{C}(G)$ with multiplication given by

$$[G/H] \cdot [G/U] = \sum_{HgU \in H \backslash G/U} [G/(H \cap {}^g U)] \quad (5.3)$$

for all $H, U \in \mathcal{C}(G)$, where $[G/(H \cap {}^gU)] = [G/K]$ for a conjugate $K \in \mathcal{C}(G)$ of $H \cap {}^gU$ in G (see, e.g., [33, 2.1]). The identity of $\Omega(G)$ is $[G/G]$.

We regard $\Omega(G)$ as $\Omega(G, A)$ with $A = \{\epsilon_A\}$. For each $X \in G\text{-set}$, the symbol $[X]$ denotes an element $\sum_{i=1}^n [G/H_i]$ of $\Omega(G)$ if $X \simeq \dot{\cup}_{i \in [n]} G/H_i$ with $H_i \in \mathcal{C}(G)$.

Remark 5.10 The product $X_1 \times X_2$ of $X_1, X_2 \in G\text{-set}$ is their cartesian product with the componentwise action of G (cf. [8, §80A]). Let $H, U \leq G$, and let $\overline{H \backslash G/U}$ be a complete set of representatives of $H \backslash G/U$. Then there exists an isomorphism

$$(G/H) \times (G/U) \xrightarrow{\sim} \dot{\bigcup}_{g \in \overline{H \backslash G/U}} G/(H \cap {}^gU), \quad (g_1H, g_2U) \mapsto g_1h(H \cap {}^gU)$$

of G -sets, where $g_2U = g_1hgU$ with $h \in H$ and $g \in \overline{H \backslash G/U}$ (see Lemma 2.15). Hence Eq.(5.3) means that $[X_1] \cdot [X_2] = [X_1 \times X_2]$ for all $X_1, X_2 \in G\text{-set}$.

Definition 5.11 We define a ring homomorphism $\alpha : \Omega(G, A) \rightarrow \Omega(G)$ by

$$[(G/U)_\tau] \mapsto [G/U]$$

for all $(U, \tau) \in \mathcal{R}(G, A)$ and define a ring homomorphism $\iota : \Omega(G) \rightarrow \Omega(G, A)$ by

$$[G/U] \mapsto [(G/U)_{1_U}]$$

for all $U \in \mathcal{C}(G)$.

Since $\alpha \circ \iota = \text{id}_{\Omega(G)}$, the Burnside ring $\Omega(G)$ is identified with $\text{Im } \iota$. We define

$$\mathcal{U}(G) := \prod_{H \in \mathcal{C}(G)} \mathbb{Z}.$$

There exists a ring monomorphism $\phi : \Omega(G) \rightarrow \mathcal{U}(G)$ given by

$$[G/U] \mapsto (|\text{inv}_H(G/U)|)_{H \in \mathcal{C}(G)}$$

for all $U \in \mathcal{C}(G)$ (cf. [8, (80.12) Proposition]), where $\text{inv}_H(G/U)$ is given by Eq.(4.1).

The ring homomorphism $\varepsilon : \mathbb{Z}H^1(H, A) \rightarrow \mathbb{Z}$ with $H \leq G$ given by

$$\sum_{\bar{\sigma} \in H^1(H, A)} \ell_{\bar{\sigma}} \bar{\sigma} \mapsto \sum_{\bar{\sigma} \in H^1(H, A)} \ell_{\bar{\sigma}}$$

for all $\ell_{\bar{\sigma}} \in \mathbb{Z}$ with $\sigma \in Z^1(H, A)$ is called the augmentation map of $\mathbb{Z}H^1(H, A)$ (cf. [21, Definition 3.2.9]).

Definition 5.12 We define a ring homomorphism $\tilde{\alpha} : \mathcal{U}(G, A) \rightarrow \mathcal{U}(G)$ by

$$(x_H)_{H \leq G} \mapsto (\varepsilon(x_H))_{H \in C(G)}$$

for all $(x_H)_{H \leq G} \in \mathcal{U}(G, A)$ and define a ring homomorphism $\tilde{\iota} : \mathcal{U}(G) \rightarrow \mathcal{U}(G, A)$ by

$$(y_H)_{H \in C(G)} \mapsto (\tilde{y}_H)_{H \leq G},$$

where $\tilde{y}_H = y_K$ for a conjugate $K \in C(G)$ of H in G , for all $(y_H)_{H \in C(G)} \in \mathcal{U}(G)$.

Obviously, $\tilde{\alpha} \circ \tilde{\iota} = \text{id}_{\mathcal{U}(G)}$. We provide the following two lemmas.

Lemma 5.13 (a) *The diagrams*

$$\begin{array}{ccc} \Omega(G, A) & \xrightarrow{\rho} & \mathcal{U}(G, A) \\ \alpha \downarrow & & \downarrow \tilde{\alpha} \\ \Omega(G) & \xrightarrow[\phi]{} & \mathcal{U}(G) \end{array} \quad \text{and} \quad \begin{array}{ccc} \Omega(G, A) & \xrightarrow{\rho} & \mathcal{U}(G, A) \\ \iota \uparrow & & \uparrow \tilde{\iota} \\ \Omega(G) & \xrightarrow[\phi]{} & \mathcal{U}(G) \end{array}$$

are commutative.

(b) *Let $x \in \Omega(G, A)$. If $\rho(x) = \tilde{\iota}(y)$ for some $y \in \mathcal{U}(G)$, then $\iota \circ \alpha(x) = x$.*

Proof. The statement (a) is clear. We prove the statement (b). Since $\tilde{\alpha} \circ \tilde{\iota} = \text{id}_{\mathcal{U}(G)}$, it follows from the statement (a) that

$$\rho \circ \iota \circ \alpha(x) = \tilde{\iota} \circ \phi \circ \alpha(x) = \tilde{\iota} \circ \tilde{\alpha} \circ \rho(x) = \tilde{\iota} \circ \tilde{\alpha} \circ \tilde{\iota}(y) = \tilde{\iota}(y) = \rho(x).$$

This, combined with Corollary 4.10, shows that $\iota \circ \alpha(x) = x$, completing the proof.

□

Lemma 5.14 (a) $\alpha \circ \eta \circ \tilde{\iota} \circ \phi = |G| \text{id}_{\Omega(G)}$. (b) $\phi \circ \alpha \circ \eta \circ \tilde{\iota} = |G| \text{id}_{\mathcal{U}(G)}$.

Proof. The lemma follows from Proposition 4.9 and Lemma 5.13(a). □

The rest of this section is devoted to the idempotents of $\mathbb{Q} \otimes_{\mathbb{Z}} \Omega(G)$.

Definition 5.15 Given $U \leq G$, we define $W_G(U)$ to be the factor group $N_G(U)/U$.

Let p be a prime or the symbol ∞ . For each $U \leq G$, we denote by $W_G(U)_p$ a Sylow p -subgroup of $W_G(U)$ provided p is a prime, and set $W_G(U)_{\infty} = W_G(U)$.

The elements $[G/H]$ for $H \in C(G)$ form a free $\mathbb{Z}_{(p)}$ -basis of the $\mathbb{Z}_{(p)}$ -lattice $\Omega(G)_{(p)}$. We identify $\mathcal{U}(G)_{(p)}$ with $\prod_{H \in C(G)} \mathbb{Z}_{(p)}$. Let $\phi^{(p)}$ denote the ring monomorphism from $\Omega(G)_{(p)}$ to $\mathcal{U}(G)_{(p)}$ determined by ϕ .

We quote [9, Proposition 1.3.5] (see also [32, Lemma 2.1]).

Proposition 5.16 *Let $\tilde{x} = (x_H)_{H \in C(G)} \in \mathcal{U}(G)_{(p)}$. Then $\tilde{x} \in \text{Im}\phi^{(p)}$ if and only if*

$$\sum_{gU \in W_G(U)_p} x_{\langle g \rangle U} \equiv 0 \pmod{|W_G(U)_p|},$$

where $x_{\langle g \rangle U} = x_K$ for a conjugate $K \in C(G)$ of $\langle g \rangle U$ in G , for all $U \in C(G)$.

Proof. The assertion follows from Theorem 5.9 and Lemma 5.13(a). $\square$

By Lemma 5.14, the primitive idempotents of $\mathbb{Q} \otimes_{\mathbb{Z}} \Omega(G)$ are the elements

$$e_H := \frac{1}{|G|} \alpha \circ \eta \circ \tilde{\iota}((\delta_H K)_{K \in C(G)}) = \frac{1}{|N_G(H)|} \sum_{U \leq H} |U| \mu(U, H) [G/U] \quad (5.4)$$

for $H \in C(G)$. This fact was shown by Gluck [14] and independently by Yoshida [31]. Obviously, $e_H e_K = \delta_{H K} e_H$ for all $H, K \in C(G)$, and $[G/G] = \sum_{H \in C(G)} e_H$.

Following [33], we present the primitive idempotents of $\Omega(G)$. Let $\sim_p$ be the equivalence relation on the set $\{(H) \mid H \leq G\}$, where (H) is the set of conjugates of H in G , generated by

$$(\langle g \rangle U) \sim_p (U)$$

for $U \leq G$ and $gU \in W_G(U)_p$ with $g \in N_G(U)$. We define an equivalence relation $\sim_p$ on the set $S(G)$ of subgroups of G by

$$H \sim_p K : \Longleftrightarrow (H) \sim_p (K).$$

Let $H \leq G$. When p is a prime, we denote by $O^p(H)$ the smallest normal subgroup of H such that $H/O^p(H)$ is a p -group (cf. [31]). Suppose that

$$H = H^{(0)} \geq H^{(1)} \geq H^{(2)} \geq \dots \geq H^{(i)} \geq \dots$$

is the derived series of H (cf. [26, Chapter 2, Definition 3.11]). Then we define $O^\infty(H) := \bigcap_{i=0}^\infty H^{(i)}$. The following lemma is well-known (cf. [33, p. 535]).

Lemma 5.17 *Let $H, U \leq G$. Then $H \sim_p U$ if and only if $(O^p(H)) = (O^p(U))$.*

Proof. The ‘if’ part follows from [26, Chapter 2, Theorem 1.6]. To prove the ‘only if’ part, we may assume that $H = \langle g \rangle U$ for some $gU \in W_G(U)_p$ with $g \in N_G(U)$. If p is a prime, then $U \geq O^p(U) \geq O^p(H)$, and hence $O^p(U) = O^p(H)$. Suppose that $p = \infty$. We have $U^{(i-1)} \geq H^{(i)} \geq U^{(i)}$ for any $i \geq 1$. If $U^{(i-1)} = U^{(i)}$ for some i , then $U^{(i-1)} = H^{(i)} = U^{(i)}$. Thus we have $O^\infty(H) = O^\infty(U)$, completing the proof. $\square$

A subgroup H of G is said to be p -perfect if $H = O^p(H)$. For each $K \leq G$, $K \sim_p O^p(K)$ by Lemma 5.17, and $O^p(K)$ is p -perfect. Let $C^{(p)}(G)$ be a full set of non-conjugate p -perfect subgroups of G . For each $H \in C^{(p)}(G)$, we define

$$e_H^{(p)} := \sum_{H \sim_p K \in C^{(p)}(G)} e_K,$$

where the sum is taken over all $K \in \mathcal{C}(G)$ such that $H \sim_p K$.

The following theorem concerns [2, Theorem 7.3] and [33, 4.12 Theorem] (see also [14, Lemma 2] and [31, Theorem 3.1]).

Theorem 5.18 *The elements $e_H^{(p)}$ for $H \in \mathcal{C}^{(p)}(G)$ are the primitive idempotents of $\Omega(G)_{(p)}$, and the elements $e_H^{(\infty)}$ for $H \in \mathcal{C}^{(\infty)}(G)$ are also those of $\Omega(G, A)$.*

Proof. For any idempotent $(x_H)_{H \in \mathcal{C}(G)}$ of $\mathcal{U}(G)_{(p)}$, it follows from Proposition 5.16 that $(x_H)_{H \in \mathcal{C}(G)} \in \text{Im} \phi^{(p)}$ if and only if $x_K = x_U \in \{0, 1\}$ for all pairs (K, U) of $K, U \in \mathcal{C}(G)$ with $K \sim_p U$. Hence the elements $e_H^{(p)}$ for $H \in \mathcal{C}^{(p)}(G)$ are the primitive idempotents of $\Omega(G)_{(p)}$. Let x be an idempotent of $\Omega(G, A)$. According to [21, Corollary 7.2.4], $\mathbb{Z}H^1(H, A)$ with $H \leq G$ contains only trivial idempotents, whence $\rho(x) = \tilde{\iota}(y)$ for some $y \in \mathcal{U}(G)$. This, combined with Lemma 5.13(b), shows that $\iota \circ \alpha(x) = x$. By this fact, we may identify x with $\alpha(x) \in \Omega(G)$. Since the map $\alpha : \Omega(G, A) \rightarrow \Omega(G)$ is a ring homomorphism, it follows that $\alpha(x)$ is an idempotent of $\Omega(G)$. Consequently, the idempotents of $\Omega(G, A)$ are those of $\Omega(G)$. This completes the proof. $\square$

There is an immediate consequence of Theorem 4.18 (see [4, 1.5. Proposition]).

Proposition 5.19 *The $\mathbb{Z}$ -module $\Omega(G, A)^\times$ has a structure of an $\Omega(G)$ -module, namely,*

$$\Omega(G) \otimes_{\mathbb{Z}} \Omega(G, A)^\times \rightarrow \Omega(G, A)^\times, \quad [G/H] \otimes_{\mathbb{Z}} x \mapsto \overline{\text{Map}}_H(G, \text{res}_H^G(x)).$$

Moreover,

$$\Omega(G, A)^\times = \prod_{H \in \mathcal{C}^{(\infty)}(G)} \{e_H^{(\infty)} x \mid x \in \Omega(G, A)^\times\},$$

where $e_H^{(\infty)} x$ denotes the effect of $e_H^{(\infty)}$ on x .

6 Units of Burnside rings

6A The Yoshida criterion for the units of Burnside rings

We turn to the unit group $\Omega(G)^\times$ of $\Omega(G)$. Let $\mathcal{U}(G)^\times$ be the unit group of $\mathcal{U}(G)$, and let $\phi^\times : \Omega(G)^\times \rightarrow \mathcal{U}(G)^\times$ be the map obtained by restriction of $\phi : \Omega(G) \rightarrow \mathcal{U}(G)$ from $\Omega(G)$ to $\Omega(G)^\times$. Obviously, $\mathcal{U}(G)^\times = \prod_{H \in \mathcal{C}(G)} \langle -1 \rangle$, where $\langle -1 \rangle = \{\pm 1\}$, and hence $\Omega(G)^\times$ is embedded in $\prod_{H \in \mathcal{C}(G)} \langle -1 \rangle$. In particular, $\Omega(G)^\times$ is an elementary abelian 2-group with identity $[G/G]$ (cf. [11, Proposition 3.1]). Thus $\Omega(G)^\times$ consists of all $x \in \Omega(G)$ such that $([G/G] \pm x)/2$ are idempotents of $\mathbb{Q} \otimes_{\mathbb{Z}} \Omega(G)$.

Example 6.1 Suppose that $K \leq G$ and $|G : K| = 2$. Then $[G/K] \cdot [G/K] = 2[G/K]$, and hence $[G/G] - [G/K] \in \Omega(G)^\times$. We have $\phi^\times([G/G] - [G/K]) = ((-1)^{\zeta(H,K)})_{H \in C(G)}$, where $\zeta(H, K) = 1$ if $H \leq K$, and $\zeta(H, K) = 0$ otherwise.

Remark 6.2 According to Dress [10], G is solvable if and only if 0 and $[G/G]$ are the only idempotents of $\Omega(G)$ (see also Lemma 5.17 and Theorem 5.18). Suppose that G is of odd order. Then by Eq.(5.4), $\Omega(G)^\times$ consists of all $x \in \Omega(G)$ such that $([G/G] \pm x)/2$ are idempotents of $\Omega(G)$, whence $|\Omega(G)^\times|$ is the number of idempotents of $\Omega(G)$. Consequently, we have $\Omega(G)^\times = \langle -[G/G] \rangle$ because, by Feit-Thompson's theorem, G is solvable (cf. [9, Proposition 1.5.1]).

Definition 6.3 Given $\tilde{x} = (x_H)_{H \in C(G)} \in \mathcal{U}(G)^\times$ and $U \leq G$, we define a class function $\gamma_U^{\tilde{x}} : W_G(U) \rightarrow \langle -1 \rangle$ by

$$gU \mapsto x_U x_{\langle g \rangle U}$$

for all $g \in N_G(U)$, where $x_{\langle g \rangle U} = x_K$ for a conjugate $K \in C(G)$ of $\langle g \rangle U$ in G .

We quote [32, Proposition 6.5] which is due to Yoshida.

Theorem 6.4 (The Yoshida criterion) *The subgroup $\text{Im } \phi^\times$ of $\mathcal{U}(G)^\times$ consists of all $\tilde{x} = (x_H)_{H \in C(G)} \in \mathcal{U}(G)^\times$ such that $\gamma_U^{\tilde{x}} \in \text{Hom}(W_G(U), \langle -1 \rangle)$ for each $U \leq G$.*

Example 6.5 Let p be an odd prime, and suppose that G is a finite p -group. Let $\tilde{x} = (x_H)_{H \in C(G)} \in \text{Im } \phi^\times$. If $G_r < \cdots < G_1 < G_0 = G$ is a sequence of subgroups of G with $|G_{i-1} : G_i| = p$ for all $i \in [r]$, then by Theorem 6.4, $x_{G_0} = x_{G_1} = \cdots = x_{G_r}$. Thus it follows from [26, Chapter 2, Theorem 1.9] that $\tilde{x}$ is determined by x_G , and hence $\tilde{x} \in \langle (-1, -1, \dots, -1) \rangle$. Consequently, $\Omega(G)^\times = \langle -[G/G] \rangle$ (see Remark 6.2).

Definition 6.6 For each $\tilde{x} \in \text{Im } \phi$, $\phi^{-1}(\tilde{x})$ denotes the unique element x of $\Omega(G)$ such that $\tilde{x} = \phi(x)$. We define a subgroup $\Omega(G)_0^\times$ of $\Omega(G)^\times$ to be the product of the subgroups $\langle [G/K] - [G/G] \rangle$ for $K \leq G$ with $|G : K| = 2$, and define a subgroup $\Omega(G)_1^\times$ of $\Omega(G)^\times$ to be the group consisting of all $x = \phi^{-1}((x_H)_{H \in C(G)})$ with $(x_H)_{H \in C(G)} \in \text{Im } \phi^\times$ such that $x_H = 1$ whenever H is cyclic.

The group $\text{Hom}(G, \langle -1 \rangle)$ with pointwise product is isomorphic to the factor group G/G_2 where G_2 is the intersection of all subgroups of index 2 in G .

Proposition 6.7 (a) $|\langle -[G/G] \rangle \times \Omega(G)_0^\times| = 2^{|\text{Hom}(G, \langle -1 \rangle)|}$.

(b) $\Omega(G)^\times = \langle -[G/G] \rangle \times \Omega(G)_0^\times \Omega(G)_1^\times \simeq \langle -[G/G] \rangle \times \text{Hom}(G, \langle -1 \rangle) \times \Omega(G)_1^\times$.

Proof. Obviously, $\Omega(G)_0^\times$ is the direct product of the subgroups $\langle [G/K] - [G/G] \rangle$ for $K \leq G$ with $|G : K| = 2$. Thus the assertion (a) holds. We prove the assertion (b). For each $K \leq G$ with $|G : K| = 2$, if $\phi([G/K] - [G/G]) = \tilde{x} = (x_H)_{H \in C(G)}$, then

by Example 6.1 and Theorem 6.4, $\gamma_{(K)} := \gamma_{\{\epsilon\}}^{\tilde{x}} \in \text{Hom}(G, \langle -1 \rangle)$, $\text{Ker } \gamma_{(K)} = K$, and $\gamma_{(K)}(g) = x_{\langle g \rangle}$ for all $g \in G$. Let $y \in \Omega(G)^\times$, and suppose that the $\{\epsilon\}$ -component of $\phi(y)$ is 1. If $\phi^\times(y) = \tilde{y} = (y_H)_{H \in \mathcal{C}(G)}$ with $y_{\{\epsilon\}} = 1$, then $\gamma_{\{\epsilon\}}^{\tilde{y}} \in \text{Hom}(G, \langle -1 \rangle)$ by Theorem 6.4, and $\gamma_{\{\epsilon\}}^{\tilde{y}}(g) = y_{\langle g \rangle}$ for all $g \in G$. This, combined with the preceding argument, shows that $([G/K] - [G/G]) \cdot y \in \Omega(G)_1^\times$ with $K = \text{Ker } \gamma_{\{\epsilon\}}^{\tilde{y}}$, which yields $y \in \Omega(G)_0^\times \Omega(G)_1^\times$. Hence $\Omega(G)_0^\times \Omega(G)_1^\times$ consists of all $x \in \Omega(G)^\times$ such that the $\{\epsilon\}$ -component of $\phi(x)$ is 1. We now obtain

$$\Omega(G)^\times = \langle -[G/G] \rangle \times \Omega(G)_0^\times \Omega(G)_1^\times.$$

Let $K_1, K_2, \dots, K_n$ be the subgroups of index 2 in G . Then $\Omega(G)_0^\times$ is the direct product of the subgroups $\langle [G/K_i] - [G/G] \rangle$ for $i \in [n]$ and $\text{Hom}(G, \langle -1 \rangle)$ is the group consisting of 1_G and the linear $\mathbb{C}$ -characters $\gamma_{(K_i)}$ for $i \in [n]$. Define a group epimorphism $\gamma : \Omega(G)_0^\times \rightarrow \text{Hom}(G, \langle -1 \rangle)$ by

$$\prod_{j=1}^m ([G/K_{i_j}] - [G/G]) \mapsto \prod_{j=1}^m \gamma_{(K_{i_j})}$$

for all sequences $(i_1, i_2, \dots, i_m)$ with $1 \leq i_1 < i_2 < \dots < i_m \leq n$ of natural numbers. Then it is obvious that $\text{Ker } \gamma = \Omega(G)_0^\times \cap \Omega(G)_1^\times$. Consequently, we have

$$\Omega(G)_0^\times \Omega(G)_1^\times \simeq \text{Hom}(G, \langle -1 \rangle) \times \Omega(G)_1^\times,$$

completing the proof. $\square$

Proposition 6.8 *Let $\widehat{\mathcal{C}}(G)$ be the set of all $U \in \mathcal{C}(G)$ such that $|N_G(U) : U| \leq 2$. For any $\tilde{x} = (x_H)_{H \in \mathcal{C}(G)} \in \text{Im } \phi^\times$, the values x_H for $H \in \mathcal{C}(G)$ are determined by the values x_U for $U \in \widehat{\mathcal{C}}(G)$. In particular, $|\Omega(G)^\times| \leq 2^{|\widehat{\mathcal{C}}(G)|}$.*

Proof. Let $\tilde{x} = (x_H)_{H \in \mathcal{C}(G)} \in \text{Im } \phi^\times$, and let $H \leq G$. By Theorem 6.4, we have

$$x_{\langle g_1 \rangle H} x_{\langle g_2 \rangle H} x_H = x_{\langle g_1 g_2 \rangle H}$$

for all $g_1, g_2 \in N_G(H)$. Hence, if $|N_G(H) : H| > 2$, then the value x_H is determined by the values x_K with $H < K \leq N_G(H)$ (cf. [7, p. 904]). This completes the proof. $\square$

Example 6.9 Assume that G is abelian. Then by Propositions 6.7 and 6.8, we have $|\Omega(G)^\times| = 2^{|\text{Hom}(G, \langle -1 \rangle)|}$, because $\widehat{\mathcal{C}}(G)$ is the set of all $K \leq G$ such that $|G : K| \leq 2$ (cf. [32, Lemma 7.1]). This fact is due to Matsuda (cf. [18, Example 4.5]).

6B Structure of the unit groups of Burnside rings

We continue to discuss the structure of $\Omega(G)^\times$.

Definition 6.10 We define a subset $\overline{C}(G)$ of $C(G)$ to be the set consisting of all subgroups U which satisfy the following conditions.

- (i) $|N_G(U) : U| \leq 2$.
- (ii) If L is a normal subgroup of U and if U/L is a non-trivial cyclic group, then U/L is a cyclic 2-group and there exists a subgroup K of index 2 in $N_G(L)$ containing L such that

$$\{\langle g \rangle L \mid g \in N_G(L) - K\} = \{\langle g \rangle L \mid g \in N_G(L) \text{ and } (\langle g \rangle L) = (U)\}.$$

Proposition 6.11 Let $U \in C(G)$, and set $\tilde{x} = ((-1)^{\delta_U H})_{H \in C(G)} \in \mathcal{U}(G)^\times$. Then $\tilde{x} \in \text{Im} \phi^\times$ if and only if $U \in \overline{C}(G)$. In particular, if $U \in \overline{C}(G)$, then $2e_U \in \Omega(G)$, or equivalently, $[G/G] - 2e_U = \phi^{-1}(\tilde{x}) \in \Omega(G)^\times$.

Proof. Assume that $\tilde{x} \in \text{Im} \phi^\times$. For any $L \leq G$, it follows from Theorem 6.4 that the map $\gamma_L^{\tilde{x}} : W_G(L) \rightarrow \langle -1 \rangle$ is a linear $\mathbb{C}$ -character of $W_G(L)$. Moreover, by assumption, $\gamma_U^{\tilde{x}}(gU) = -1$ for any $g \in N_G(U) - U$. This means that $\text{Ker } \gamma_U^{\tilde{x}} = U/U$. Consequently, $|N_G(U) : U| \leq 2$. Let L be a normal subgroup of U , and suppose that U/L is non-trivial cyclic. Set $U = \langle r \rangle L$ with $r \in N_G(L) - L$. Then for any $g \in N_G(L)$, $\gamma_L^{\tilde{x}}(gL) = -1$ if and only if $\langle g \rangle L$ is a conjugate of $\langle r \rangle L$ in G . In particular, rL must be a 2-element of $W_G(L)$, whence U/L is a cyclic 2-group. Moreover, there exists a subgroup K of index 2 in $N_G(L)$ containing L such that $K/L = \text{Ker } \gamma_L^{\tilde{x}}$ and

$$\{\langle g \rangle L \mid g \in N_G(L) - K\} = \{\langle g \rangle L \mid g \in N_G(L) \text{ and } (\langle g \rangle L) = (U)\}.$$

Thus $U \in \overline{C}(G)$, as required. Conversely, if $U \in \overline{C}(G)$, then by Theorem 6.4, we have $\tilde{x} \in \text{Im} \phi^\times$, completing the proof. $\square$

Remark 6.12 Under the hypotheses of Proposition 6.11, it follows from Eq.(5.4) that $\tilde{x} \in \text{Im} \phi^\times$ if and only if $[G/G] - 2e_U \in \Omega(G)^\times$.

Corollary 6.13 Let $U \in \overline{C}(G)$, and suppose that U is non-trivial cyclic. Then U is a Sylow 2-subgroup of G , and $N_G(U) = U$.

Proof. Set $\tilde{x} = ((-1)^{\delta_U H})_{H \in C(G)} \in \mathcal{U}(G)^\times$. By Theorem 6.4 and Proposition 6.11, the map $\gamma_{\{\epsilon\}}^{\tilde{x}} : G \rightarrow \langle -1 \rangle$ is a linear $\mathbb{C}$ -character of G . Since U is non-trivial cyclic, it follows that $\gamma_{\{\epsilon\}}^{\tilde{x}}$ is not the trivial character of G . If $K = \text{Ker } \gamma_{\{\epsilon\}}^{\tilde{x}}$, then any cyclic subgroup $\langle g \rangle$ with $g \in G - K$ is a conjugate of U in G and

$$\frac{|G|}{2} = |K| = |G - K| = |G : N_G(U)| \cdot \frac{|U|}{2} = \frac{|G|}{2|N_G(U) : U|}$$

because U is a 2-group. Thus we have $|N_G(U) : U| = 1$. The corollary is now a consequence of [26, Chapter 2, Theorem 1.6]. This completes the proof. $\square$

Let $\lambda = (\lambda_1, \dots, \lambda_j, \dots, \lambda_m, \lambda_{m+1}, \dots)$, where $\lambda_1 > \dots > \lambda_j > \dots > \lambda_m > 0$ and $\lambda_\ell = 0$ for $\ell = m+1, m+2, \dots$, be a partition of $n \in \mathbb{N}$. Such a partition is said to be strict. We set $S_\lambda = S_{(\lambda_1)} \times \dots \times S_{(\lambda_j)} \times \dots \times S_{(\lambda_m)}$, where each $S_{(\lambda_j)}$ is the symmetric group on $\{\sum_{i \geq j+1} \lambda_i + 1, \dots, \sum_{i \geq j} \lambda_i\}$. Let S_n be the symmetric group on $[n]$. Then S_λ is a Young subgroup of S_n associated with the strict partition λ .

Proposition 6.14 *For any strict partition λ of n , the set $\overline{C}(S_n)$ contains a conjugate of the Young subgroup S_λ of S_n associated with λ .*

Proof. We may assume that $S_\lambda \in C(S_n)$. Obviously, $N_{S_n}(S_\lambda) = S_\lambda$. We show that $S_\lambda \in \overline{C}(S_n)$. Under the preceding notation, let $A_{(\lambda_j)}$ with $j \in [m]$ be the subgroup of $S_{(\lambda_j)}$ consisting of all even permutations. Then the commutator subgroup of S_λ is $A_{(\lambda_1)} \times \dots \times A_{(\lambda_j)} \times \dots \times A_{(\lambda_m)}$. Hence every normal subgroup L of S_λ such that S_λ/L is non-trivial cyclic is a subgroup of index 2 in S_λ . If $N_{S_n}(L) = S_\lambda$ for a subgroup L of index 2 in S_λ , then $\langle g \rangle L = S_\lambda$ for any $g \in N_{S_n}(L) - L$. Thus it suffices to verify that, if $N_{S_n}(L) \neq S_\lambda$ for a subgroup L of index 2 in S_λ , then

$$\{\langle g \rangle L \mid g \in N_{S_n}(L) - K\} = \{\langle g \rangle L \mid g \in N_{S_n}(L) \text{ and } (\langle g \rangle L) = (S_\lambda)\}$$

for a subgroup K of index 2 in $N_{S_n}(L)$ containing L . Let $L \leq S_\lambda$ with $|S_\lambda : L| = 2$ and $N_{S_n}(L) \neq S_\lambda$. Then $\lambda_{m-1} = 2$, $\lambda_m = 1$, and every permutation in L fixes both $2 \in [n]$ and $3 \in [n]$. (In this case, $S_{(\lambda_{m-1})}$ is the symmetric group on $\{2, 3\}$). Hence it turns out that $L = S_{(\lambda_1)} \times \dots \times S_{(\lambda_j)} \times \dots \times S_{(\lambda_{m-2})}$, $S_\lambda = L \times S_{(\lambda_{m-1})} \times S_{(\lambda_m)}$, and $N_{S_n}(L) = L \times S_3$. Consequently, $L \leq L \times A_3 \leq N_{S_n}(L)$, $|N_{S_n}(L) : L \times A_3| = 2$, $(\langle g \rangle L) \neq (S_\lambda)$ for any $g \in L \times A_3$, where A_3 is the alternating group on $[3]$, and the set of conjugates of S_λ in S_n includes the set $\{\langle g \rangle L \mid g \in N_{S_n}(L) - (L \times A_3)\}$, as required. We now conclude that $S_\lambda \in \overline{C}(S_n)$, completing the proof. $\square$

Definition 6.15 For each $L \leq G$, we define a subset $S(G; L)$ of $S(G)$ to be the set consisting of all subgroups U of $N_G(L)$ which satisfy the following conditions.

- (i) U/L is a non-trivial cyclic 2-group.
- (ii) There exists a subgroup K of index 2 in $N_G(L)$ containing L such that

$$\{\langle g \rangle L \mid g \in N_G(L) - K\} = \{\langle g \rangle L \mid g \in N_G(L) \text{ and } (\langle g \rangle L) = (U)\}.$$

Let $\approx$ be the equivalence relation on the set $\{(H) \mid G \geq H \neq \{\epsilon\}\}$ generated by

$$(\langle g \rangle L) \approx (L)$$

for $L \in C(G)$ and $g \in N_G(L)$ such that $\langle g \rangle L \notin S(G; L)$. We set $C(G)^\circ = C(G) - \{\epsilon\}$, and define an equivalence relation $\approx$ on $C(G)^\circ$ by

$$H \approx K : \Longleftrightarrow (H) \approx (K).$$

Proposition 6.16 *If $|G| > 2$, then each $U \in \overline{C}(G)$ forms an equivalence class consisting of a single element with respect to the equivalence relation $\approx$ on $C(G)^\circ$.*

Proof. Suppose that $|G| > 2$, and let $U \in \overline{C}(G)$. Then $U \neq \{\epsilon\}$ and $|N_G(U) : U| \leq 2$. If $N_G(U) \neq U$, then $|N_G(U) : U| = 2$ and $N_G(U) \in S(G; U)$. Moreover, if L is a normal subgroup of U and if U/L is a non-trivial cyclic group, then $U \in S(G; L)$. Thus (U) is isolated with respect to $\approx$. This completes the proof. $\square$

Proposition 6.17 *Suppose that $\tilde{y} = (y_H)_{H \in C(G)} \in \text{Im } \phi^\times$ and $\phi^{-1}(\tilde{y}) \in \Omega(G)_1^\times$. Let $U \in C(G)^\circ$, and define $\tilde{x} = (x_H)_{H \in C(G)} \in \mathcal{U}(G)^\times$ by*

$$x_H = \begin{cases} y_H & \text{if } H \approx U, \\ 1 & \text{if } H \not\approx U \text{ or } H = \{\epsilon\}. \end{cases}$$

Then $\tilde{x} \in \text{Im } \phi^\times$ and $\phi^{-1}(\tilde{x}) \in \Omega(G)_1^\times$.

Proof. By the definition of $\tilde{x}$, the map $\gamma_{\{\epsilon\}}^{\tilde{x}} : G \rightarrow \langle -1 \rangle$ is the trivial character of G . Hence it suffices to verify that $\tilde{x} \in \text{Im } \phi^\times$. Let $L \in C(G)^\circ$. We show that the map $\gamma_L^{\tilde{x}} : W_G(L) \rightarrow \langle -1 \rangle$ is a linear $\mathbb{C}$ -character of $W_G(L)$. By Theorem 6.4, the map $\gamma_L^{\tilde{y}} : W_G(L) \rightarrow \langle -1 \rangle$ is a linear $\mathbb{C}$ -character of $W_G(L)$. We may assume that $\gamma_L^{\tilde{x}} \notin \{\gamma_L^{\tilde{y}}, 1_{W_G(L)}\}$. (If $\langle g \rangle L \notin S(G; L)$ for all $g \in N_G(L) - L$, then either $\gamma_L^{\tilde{x}} = \gamma_L^{\tilde{y}}$ or $\gamma_L^{\tilde{x}} = 1_{W_G(L)}$.) Obviously, $\gamma_L^{\tilde{x}}(L) = 1$. We analysis the values $\gamma_L^{\tilde{x}}(\langle g \rangle L)$ for $g \in N_G(L) - L$ in each of the cases where $L \approx U$ and $L \not\approx U$. Let r be any element of $N_G(L) - L$ such that $\langle r \rangle L \in S(G; L)$. Then there exist a subgroup K of index 2 in $N_G(L)$ containing L such that for each $g \in N_G(L)$, $g \in N_G(L) - K$ if and only if $\langle g \rangle L$ is a conjugate of $\langle r \rangle L$ in G . We define a map $\beta_r : W_G(L) \rightarrow \langle -1 \rangle$ to be the linear $\mathbb{C}$ -character of $W_G(L)$ whose kernel is K/L .

Case 1. Assume that $L \approx U$. Let $\mathcal{X} = \{\langle r_i \rangle L \mid i \in [\ell]\}$ be a full set of non-conjugate subgroups of G chosen from among the subgroups $\langle g \rangle L$ for $g \in N_G(L) - L$ with $\gamma_L^{\tilde{x}}(gL) \neq \gamma_L^{\tilde{y}}(gL)$. Then we have $(\langle r_i \rangle L) \not\approx (L)$ and $\langle r_i \rangle L \in S(G; L)$ for all $i \in [\ell]$. For any $g \in N_G(L) - L$,

$$\gamma_L^{\tilde{x}}(gL) = -\gamma_L^{\tilde{y}}(gL) = \gamma_L^{\tilde{y}}(gL) \prod_{i=1}^{\ell} \beta_{r_i}(gL)$$

if $\langle g \rangle L$ is a conjugate of some $\langle r_j \rangle L$ with $j \in [\ell]$ in G , and

$$\gamma_L^{\tilde{x}}(gL) = \gamma_L^{\tilde{y}}(gL) = \gamma_L^{\tilde{y}}(gL) \prod_{i=1}^{\ell} \beta_{r_i}(gL)$$

otherwise. Thus we have

$$\gamma_L^{\tilde{x}} = \gamma_L^{\tilde{y}} \prod_{i=1}^{\ell} \beta_{r_i}.$$

Case 2. Assume that $L \not\approx U$. Then $x_L = 1$. Let $\mathcal{Y} = \{\langle r_i \rangle L \mid i \in [\ell]\}$ be a full set of non-conjugate subgroups of G chosen from among the subgroups $\langle g \rangle L$ for $g \in N_G(L) - L$ with $\gamma_L^{\tilde{x}}(gL) \neq 1$. Then $(\langle r_i \rangle L) \approx (U)$, whence $(\langle r_i \rangle L) \not\approx (L)$ and $\langle r_i \rangle L \in S(G; L)$ for all $i \in [\ell]$. By an argument analogous to that in Case 1, we have

$$\gamma_L^{\tilde{x}} = \prod_{i=1}^{\ell} \beta_{r_i}.$$

We now conclude that the map $\gamma_L^{\tilde{x}} : W_G(L) \rightarrow \langle -1 \rangle$ is a linear $\mathbb{C}$ -character of $W_G(L)$ in either case. Consequently, $\gamma_L^{\tilde{x}} \in \text{Hom}(W_G(L), \langle -1 \rangle)$ for any $L \leq G$. This, combined with Theorem 6.4, shows that $\tilde{x} \in \text{Im} \phi^\times$, completing the proof. $\square$

Corollary 6.18 *Let $C(G)^\circ / \approx$ be a complete set of representatives of equivalence classes with respect to the equivalence relation $\approx$ on $C(G)^\circ$. Set*

$$\Omega(G)_U^\times = \{\phi^{-1}(\tilde{x}) \mid \tilde{x} = (x_H)_{H \in C(G)} \in \text{Im} \phi^\times \text{ and } x_H = 1 \text{ if } H \not\approx U \text{ or } H = \{\epsilon\}\}$$

for each $U \in C(G)^\circ / \approx$. Then

$$\Omega(G)_1^\times = \prod_{U \in C(G)^\circ / \approx} (\Omega(G)_U^\times \cap \Omega(G)_1^\times).$$

Moreover, if $U \in \overline{C}(G) \cap C(G)^\circ$, then $U \in C(G)^\circ / \approx$ and $\Omega(G)_U^\times = \langle [G/G] - 2e_U \rangle$.

Proof. The assertion follows from Propositions 6.11, 6.16, and 6.17. $\square$

7 Units of monomial Burnside rings

7A The unit groups of monomial Burnside rings

We continue assuming that A is abelian. Given a commutative unital ring R , we denote by $R^\times$ the unit group of R , and denote by R^ω the group of torsion units of R . For each $H \leq G$, since $H^1(H, A)$ is a finite abelian group, it follows from [21, Theorem 8.3.1] that $(\mathbb{Z}H^1(H, A))^\times$ is a finitely generated abelian group.

Lemma 7.1 *The group $\mathcal{U}(G, A)^\times$ is a finitely generated abelian group.*

Proof. Observe that $\mathcal{U}(G, A) \simeq \prod_{H \in C(G)} (\mathbb{Z}H^1(H, A))^{N_G(H)}$, where

$$(\mathbb{Z}H^1(H, A))^{N_G(H)} = \{x_H \in \mathbb{Z}H^1(H, A) \mid \text{con}_H^g(x_H) = x_H \text{ for all } g \in N_G(H)\}.$$

Then we have $\mathcal{U}(G, A)^\times \simeq \prod_{H \in C(G)} J_H$, where

$$J_H = (\mathbb{Z}H^1(H, A))^\times \cap (\mathbb{Z}H^1(H, A))^{N_G(H)}.$$

Hence it suffices to verify that the groups J_H for $H \leq G$ are finitely generated. Let $H \leq G$, and assume that $(\mathbb{Z}H^1(H, A))^\times$ is generated by $x_1, \dots, x_k$. We set $y_i = \prod_{g \in N_G(H)} \text{con}_H^g(x_i)$ for all i , and set $\widehat{J}_H = \langle y_1, \dots, y_k \rangle$. Obviously, $\widehat{J}_H$ is a subgroup of J_H . We have

$$x^{|N_G(H)|} = \prod_{g \in N_G(H)} \text{con}_H^g(x) \in \widehat{J}_H$$

for any $x \in J_H$, so that $J_H/\widehat{J}_H$ is a torsion subgroup of $(\mathbb{Z}H^1(H, A))^\times/\widehat{J}_H$. Since $(\mathbb{Z}H^1(H, A))^\times/\widehat{J}_H$ is finitely generated, it follows from the fundamental theorem of abelian groups (see, e.g., [16, I, §10, Theorem 8]) that $J_H/\widehat{J}_H$ is a finite group. Thus J_H is finitely generated, as desired. This completes the proof. $\square$

Proposition 7.2 *The group $\Omega(G, A)^\times$ is a finitely generated abelian group. In particular, $\Omega(G, A)^\times$ is the direct product of $\Omega(G, A)^\omega$ and a free abelian group of finite rank, and $\Omega(G, A)^\omega$ is a finite abelian group.*

Proof. By the fundamental theorem of abelian groups, it suffices to prove the first statement. Using Proposition 5.2 and Corollary 5.5, we have

$$\mathbb{Q} \otimes_{\mathbb{Z}} \text{Im} \rho = \mathbb{Q} \otimes_{\mathbb{Z}} \mathcal{U}(G, A).$$

This, combined with [21, Lemma 2.9.5], shows that $|\mathcal{U}(G, A)^\times : (\text{Im} \rho)^\times|$ is finite. Moreover, by Lemma 7.1, $\mathcal{U}(G, A)^\times$ is finitely generated. Hence it follows from [17, Corollary 2.7.1] that $(\text{Im} \rho)^\times$ is finitely generated. By Corollary 4.10, we have $\Omega(G, A)^\times \simeq (\text{Im} \rho)^\times$, completing the proof. $\square$

7B Torsion units of monomial Burnside rings

From Higman's theorem (cf. [21, Theorem 7.1.4]), we know that for any $H \leq G$,

$$(\mathbb{Z}H^1(H, A))^\omega = \langle -1 \rangle \times H^1(H, A) = \{\pm \bar{\sigma} \mid \sigma \in Z^1(H, A)\}. \quad (7.1)$$

Theorem 7.3 *The necessary and sufficient condition for an element $\tilde{x} = (x_H)_{H \leq G}$ of $\mathcal{U}(G, A)^\omega$ to be contained in $\text{Im} \rho$ is that $\gamma_U^{\tilde{\alpha}(\tilde{x})} \in \text{Hom}(W_G(U), \langle -1 \rangle)$ for all $U \leq G$ and $(\varepsilon(x_H)x_H)_{H \leq G} \in \Upsilon(G, A)$ (see Definitions 5.12 and 6.3), where*

$$\Upsilon(G, A) = \left\{ (\overline{\sigma_H})_{H \leq G} \in \mathcal{U}(G, A)^\omega \mid \begin{array}{l} \sigma_U \in Z^1(U, A) \text{ and } \overline{\sigma_U} = \text{res}_U^{(g)U}(\overline{\sigma_{\langle g \rangle U}}) \\ \text{for all } U \leq G \text{ and } g \in N_G(U) \end{array} \right\}.$$

Proof. Let $\tilde{x} = (x_H)_{H \leq G} \in \mathcal{U}(G, A)^\omega$. Suppose that for each $H \leq G$, $\overline{\sigma_H} = \varepsilon(x_H)x_H$ with $\sigma_H \in Z^1(H, A)$ (see Eq.(7.1)). We first prove 'sufficient' part. By assumption, $\text{con}_U^g(\overline{\sigma_U}) = \overline{\sigma_U}$ and $\overline{\sigma_U} = \text{res}_U^{(g)U}(\overline{\sigma_{\langle g \rangle U}})$ for all $U \leq G$ and $g \in N_G(U)$, so that

$$\psi \circ \kappa^{-1}(\tilde{x}) = (z_{(U, \tau)} \bmod |W_G(U, \tau)|)_{(U, \tau) \in \mathcal{R}(G, A)},$$

where

$$z_{(U,\tau)} = \begin{cases} \sum_{gU \in W_G(U)} \varepsilon(x_{\langle g \rangle U}) & \text{if } \bar{\tau} = \overline{\sigma_U}, \\ 0 & \text{otherwise.} \end{cases}$$

For any $U \leq G$, since $\gamma_U^{\tilde{\alpha}(\tilde{x})} \in \text{Hom}(W_G(U), \langle -1 \rangle)$, we have

$$\frac{1}{|W_G(U)|} \sum_{gU \in W_G(U)} \varepsilon(x_U) \varepsilon(x_{\langle g \rangle U}) \in \{0, 1\}$$

by [8, (9.21) Proposition]. Hence either $z_{(U,\tau)} = \varepsilon(x_U)|W_G(U)|$ or $z_{(U,\tau)} = 0$ for all $(U, \tau) \in \mathcal{R}(G, A)$, which yields $\psi \circ \kappa^{-1}(\tilde{x}) = 0 \in \text{Obs}(G, A)$. This, combined with Proposition 5.2 and Theorem 5.9, shows that $\tilde{x} \in \text{Im } \rho$, as desired. We next prove ‘necessary’ part. Assume that $\rho(x) = \tilde{x}$ with $x \in \Omega(G, A)^\omega$. Then $\alpha(x) \in \Omega(G)^\times$, because the map $\alpha : \Omega(G, A) \rightarrow \Omega(G)$ is a ring homomorphism. By Lemma 5.13(a) and Theorem 6.4, $\gamma_U^{\tilde{\alpha}(\tilde{x})} \in \text{Hom}(W_G(U), \langle -1 \rangle)$ for all $U \leq G$, and

$$\rho(\iota \circ \alpha(x) \cdot x) = (\overline{\sigma_H})_{H \leq G} \in \mathcal{U}(G, A)^\omega.$$

In particular, we have $\text{con}_U^g(\overline{\sigma_U}) = \overline{\sigma_U}$ for all $U \leq G$ and $g \in N_G(U)$. For each $(U, \tau) \in \mathcal{R}(G, A)$ with $\bar{\tau} = \overline{\sigma_U}$, the (U, τ) -component of $\psi \circ \kappa^{-1}((\overline{\sigma_H})_{H \leq G})$ is

$$\sum_{gU \in W_G(U), \overline{\sigma_U} = \text{res}_U^{\langle g \rangle U}(\overline{\sigma_{\langle g \rangle U})} 1 \bmod |W_G(U)|,$$

where the sum is taken over all left cosets gU , $g \in N_G(U)$, of U in $N_G(U)$ such that $\overline{\sigma_U} = \text{res}_U^{\langle g \rangle U}(\overline{\sigma_{\langle g \rangle U})}$. Since $(\overline{\sigma_H})_{H \leq G} \in \text{Im } \rho$, it follows from Proposition 5.2 and Theorem 5.9 that $\overline{\sigma_U} = \text{res}_U^{\langle g \rangle U}(\overline{\sigma_{\langle g \rangle U})}$ for all $U \leq G$ and $g \in N_G(U)$, as desired. This completes the proof. $\square$

In §4A, the ring epimorphism $\rho_G^G : \Omega(G, A) \rightarrow \mathbb{Z}H^1(G, A)$ is given by

$$[(G/U)_\tau] \mapsto \begin{cases} \bar{\tau} & \text{if } G = U, \\ 0 & \text{otherwise} \end{cases}$$

for all $(U, \tau) \in \mathcal{R}(G, A)$ (see Lemma 4.6(a)). Following [2, §7], we define a ring monomorphism $v : \mathbb{Z}H^1(G, A) \rightarrow \Omega(G, A)$ by

$$\bar{\chi} \mapsto [(G/G)_\chi]$$

for all $\chi \in Z^1(G, A)$ (see Lemmas 2.6 and 2.15). There are group homomorphisms

$$v^\omega : (\mathbb{Z}H^1(G, A))^\omega \rightarrow \Omega(G, A)^\omega \quad \text{and} \quad \theta^\omega : \Omega(G, A)^\omega \rightarrow (\mathbb{Z}H^1(G, A))^\omega$$

inherited from v and ρ_G^G , respectively (see Eq.(7.1)). Hence it turns out that

$$\Omega(G, A)^\omega = \text{Im } v^\omega \times \text{Ker } \theta^\omega \simeq \langle -1 \rangle \times H^1(G, A) \times \text{Ker } \theta^\omega$$

(cf. [2, §8]), because $\theta^\omega \circ v^\omega = \text{id}_{(\mathbb{Z}H^1(G, A))^\omega}$. We continue to describe $\Omega(G, A)^\omega$.

Corollary 7.4 *Identify the finite groups $\Omega(G)^\times$ and $H^1(G, A)$ with the subgroups $\{\iota(u) \mid u \in \Omega(G)^\times\}$ and $\{[(G/G)_\chi] \mid \chi \in Z^1(G, A)\}$ of $\Omega(G, A)^\omega$, respectively. Set*

$$\nabla(G, A) = \left\{ \frac{1}{|G|} \sum_{H \leq G} \sum_{U \leq H} |U| \mu(U, H) [(G/U)_{\sigma_H|U}] \mid \begin{array}{l} (\overline{\sigma_H})_{H \leq G} \in \Upsilon(G, A) \\ \text{with } \sigma_G = 1_G \end{array} \right\}.$$

Then

$$\Omega(G, A)^\omega = \Omega(G)^\times \times H^1(G, A) \times \nabla(G, A).$$

Proof. Let $x \in \Omega(G, A)^\omega$, and suppose that $\rho(x) = (x_H)_{H \leq G}$. By Theorem 7.3, $\rho(\iota \circ \alpha(x) \cdot x) = (\varepsilon(x_H)x_H)_{H \leq G} \in \Upsilon(G, A)$. Since the map $\alpha : \Omega(G, A) \rightarrow \Omega(G)$ is a ring epimorphism, it follows from Proposition 4.9 and Theorem 7.3 that

$$\Omega(G, A)^\omega = \Omega(G)^\times \times \left\{ \frac{1}{|G|} \eta((\overline{\sigma_H})_{H \leq G}) \mid (\overline{\sigma_H})_{H \leq G} \in \Upsilon(G, A) \right\}.$$

Moreover, $\rho([(G/G)_\chi]) = (\text{res}_H^G(\overline{\chi}))_{H \leq G} \in \Upsilon(G, A)$ for all $\chi \in Z^1(G, A)$, and hence

$$\Upsilon(G, A) = \{\rho([(G/G)_\chi]) \mid \chi \in Z^1(G, A)\} \times \{(\overline{\sigma_H})_{H \leq G} \in \Upsilon(G, A) \mid \sigma_G = 1_G\}.$$

The assertion now follows from Proposition 4.9. This completes the proof. $\square$

Remark 7.5 Suppose that G is of odd order and that G acts trivially on A . Then by Remark 6.2 and Corollary 7.4, we have

$$\Omega(G, A)^\omega = \langle -[(G/G)_{1_G}] \rangle \times \Omega(G, A)^{\text{odd}},$$

where $\Omega(G, A)^{\text{odd}}$ is the Hall $2'$ -subgroup of $\Omega(G, A)^\omega$ (cf. [2, Proposition 8.2]).

Example 7.6 Suppose that G is nilpotent. Then by [26, Chapter 4, Theorem 2.9], $\nabla(G, A) = \langle [(G/G)_{1_G}] \rangle$ in Corollary 7.4, and hence

$$\Omega(G, A)^\omega \simeq \Omega(G)^\times \times H^1(G, A).$$

REFERENCES

- [1] M. Aigner, Combinatorial Theory, Grundlehren der Mathematischen Wissenschaften, 234, Springer-Verlag, Berlin-New York, 1979.
- [2] L. Barker, Fibred permutation sets and the idempotents and units of monomial Burnside rings, J. Algebra **281** (2004), 535–566.
- [3] R. Boltje, A canonical Brauer induction formula, Astérisque, **181–182** (1990), 31–59.

- [4] R. Boltje, A general theory of canonical induction formulae, *J. Algebra* **206** (1998), 293–343.
- [5] S. Bouc, Burnside rings, *Handbook of algebra*, Vol. 2, 739–804, North-Holland, Amsterdam, 2000.
- [6] S. Bouc, The functor of units of Burnside ring for p -groups, *Comment. Math. Helv.* **82** (2007), 583–615.
- [7] S. Bouc, The slice Burnside ring and the section Burnside ring of a finite group, *Compositio Math.* **148** (2012), 868–906.
- [8] C. W. Curtis and I. Reiner, *Methods of Representation Theory*, Vol. I, II, Wiley-Interscience, New York, 1981, 1987.
- [9] T. tom Dieck, *Transformation Groups and Representation Theory*, Lecture Notes in Mathematics, 766, Springer, Berlin, 1979.
- [10] A. Dress, A characterisation of solvable groups, *Math. Z.* **110** (1969), 213–217.
- [11] A. Dress, Operations in representation rings, in “Representation theory of finite groups and related topics,” (Madison, Wis., 1970), 39–45, *Proc. Sympos. Pure Math.*, Vol. XXI, Amer. Math. Soc., Providence, R.I., 1971.
- [12] A. Dress, The ring of monomial representations I. Structure theory, *J. Algebra*, **18** (1971), 137–157.
- [13] B. Fotsing and B. Külshammer, Modular species and prime ideals for the ring of monomial representations of a finite group, *Comm. Algebra* **33** (2005), 3667–3677.
- [14] D. Gluck, Idempotent formula for the Burnside algebra with applications to the p -subgroup simplicial complex, *Illinois J. Math.* **25** (1981), 63–67.
- [15] H. Idei and F. Oda, The table of marks, the Kostka matrix, and the character table of the symmetric group, *J. Algebra* **429** (2015), 318–323.
- [16] S. Lang, *Algebra*, Addison-Wesley, Reading, 1965.
- [17] W. Magnus, A. Karrass, and D. Solitar, *Combinatorial Group Theory. Presentations of groups in terms of generators and relations*, Reprint of the 1976 second edition, Dover Publications, Inc., Mineola, NY, 2004.
- [18] T. Matsuda, On the unit groups of Burnside rings, *Japan. J. Math. (N.S.)* **8** (1982), 71–93.
- [19] T. Matsuda, A note on the unit groups of Burnside rings as Burnside ring modules, *J. Fac. Sci. Shinshu Univ.* **21** (1986), 1–10.

- [20] T. Matsuda and T. Miyata, On the unit groups of Burnside rings of finite groups, *J. Math. Soc. Japan* **35** (1983), 345–354.
- [21] C. P. Milies and S. K. Sehgal, *An Introduction to Group Rings*, Kluwer Academic Publishers, Dordrecht, 2002.
- [22] M. Müller, On the isomorphism problem for the ring of monomial representations of a finite group, *J. Algebra* **333** (2011), 427–457.
- [23] M. Müller, Isomorphic rings of monomial representations, *J. Algebra* **367** (2012), 105–119.
- [24] F. Oda, Y. Takegahara, and T. Yoshida, The units of a partial Burnside ring relative to the Young subgroups of a symmetric group, *J. Algebra*, **460** (2016), 370–379.
- [25] R. P. Stanley, *Enumerative Combinatorics, Vol. I*, Cambridge University Press, Cambridge, 1997.
- [26] M. Suzuki, *Group Theory I, II*, Springer-Verlag, New York, 1982, 1986.
- [27] Y. Takegahara, Multiple Burnside rings and Brauer induction formulae, *J. Algebra* **324** (2010), 1656–1686.
- [28] Y. Takegahara, Induction formulae for Mackey functors with applications to representations of the twisted quantum double of a finite group, *J. Algebra* **410** (2014), 85–147.
- [29] J. Thévenaz, Permutation representations arising from simplicial complexes, *J. Combin. Theory Ser. A* **46** (1987), 121–155.
- [30] E. Yalçın, An induction theorem for the unit groups of Burnside rings of 2-groups, *J. Algebra* **289** (2005), 105–127.
- [31] T. Yoshida, Idempotents of Burnside rings and Dress induction theorem, *J. Algebra* **80** (1983), 90–105.
- [32] T. Yoshida, On the unit groups of Burnside rings, *J. Math. Soc. Japan* **42** (1990), 31–64.
- [33] T. Yoshida, The generalized Burnside ring of a finite group, *Hokkaido Math. J.* **19** (1990), 509–574.