Zero potential energy criterion applied to Hartree Fock wave function s

著者	KOGA Toshikatsu
journal or publication title	The journal of chemical physics
volume	89
number	7
page range	4209-4211
year	1988-10-01
URL	http://hdl.handle.net/10258/932

doi: info:doi/10.1063/1.455688

Zero potential energy criterion applied to Hartree-Fock wave functions

Toshikatsu Koga

Department of Applied Chemistry and Department of Applied Science for Energy, Muroran Institute of Technology, Muroran, Hokkaido 050, Japan

(Received 1 March 1988; accepted 24 June 1988)

The zero potential energy criterion, which is a necessary condition for the exact wave function, is applied to Hartree-Fock wave functions for a closed-shell system with doubly occupied spatial orbitals. With the help of the known long-range asymptotic behavior of Hartree-Fock orbitals, we first derive a single-electron zero potential energy criterion to be satisfied by Hartree-Fock orbitals. The Hartree-Fock wave function is then shown to never satisfy the zero potential energy criterion, which implies that the Hartree-Fock approximation cannot describe the correct long-range asymptotic behavior of many-electron wave functions. Some numerical illustrations are given.

I. INTRODUCTION

Previously, the zero potential energy (E_{zp}) criterion has been proposed¹ and applied¹⁻³ as a sensitive measure for assessing the accuracy of approximate wave functions. E_{zp} is defined as the local energy⁴⁻⁶ evaluated at those points in position space at which the potential energy operator $V(\{\mathbf{r}_i\})$ vanishes, i.e.,

$$E_{zp} = \lim_{V \rightarrow 0} \{ [T(\{\mathbf{r}_i\})\Psi(\{\mathbf{r}_i\})] / \Psi(\{\mathbf{r}_i\}) \}, \quad (1)$$

where $\Psi(\{\mathbf{r}_i\})$ is the total electronic wave function, $\{\mathbf{r}_i\}$ the position vectors of electrons, and $T(\{\mathbf{r}_i\})$ the electronic kinetic energy operator $T(\{\mathbf{r}_i\}) = \sum_i t(\mathbf{r}_i)$, where $t(\mathbf{r}_i) = -(1/2)\Delta_i$. The zero potential energy has an important property¹ that the relation

$$E_{zp} = E_{av} = E \quad (2)$$

is a necessary (but not sufficient) condition for a wave function to be the true wave function, where E is the true energy and E_{av} the average energy. It is known¹ that E_{zp} is more sensitive to the wave function error than E_{av} , since E_{zp} and E_{av} are, respectively, linear and quadratic in the error.

The zero potential energy formula, Eq. (1), is a special form of the local energy formula with vanishing potential energy contribution, and it may be regarded as a partner to the zero momentum energy criterion⁷⁻¹⁰ which considers the momentum-space local energy with vanishing kinetic energy contribution. For atoms and molecules of our interest, the limiting procedure $V \rightarrow 0$ involved in Eq. (1) is equivalent to considering a point infinitely apart from the nuclei where Coulombic interactions disappear. Therefore, E_{zp} measures the quality of wave functions at their long-range tails.

In the present paper, we study the zero potential energy of the Hartree-Fock wave function for a closed-shell system from the viewpoints of the one-electron orbital and the total electronic wave function. In Sec. II, the zero potential consideration is applied to the Fock equation and the orbitalwise zero potential energy criterion is derived based on the known long-range asymptotic behavior^{11,12} (see also Refs. 13-19 and the references therein). In Sec. III, the total zero potential energy is derived for the Hartree-Fock wave function described as a single Slater determinant. The result shows that the Hartree-Fock approximation cannot correctly de-

scribe the long-range behavior of the many-electron wave function. Some numerical examples are presented in Sec. IV. Atomic units are used throughout this paper.

II. ORBITALWISE ZERO POTENTIAL ENERGY

Let us consider the Hartree-Fock approximation for a closed-shell $2N$ electron atomic or molecular system with N doubly occupied spatial orbitals $\{\psi_i\}$. For a large r , the leading asymptotic behavior of the Hartree-Fock orbital ψ_i is known^{11,12} to be

$$\psi_i(\mathbf{r}) = [a_i r^{b_i} + O(r^{b_i-1})] \exp(-\zeta_i r), \quad r = |\mathbf{r}|. \quad (3)$$

The exponent ζ_i is $(-2\epsilon_i)^{1/2}$ for atoms in which only s -type orbitals appear (referred to as s -type atoms hereafter), while otherwise it is $(-2\epsilon_h)^{1/2}$ independent of i . ϵ_i is the orbital energy associated with ψ_i , and h refers to the highest occupied orbital which is presumed to be nondegenerate. The expression for b_i can be found in Ref. 12. Now from Eq. (3), the zero potential energy criterion for the orbital ψ_i is obtained as

$$\begin{aligned} \epsilon_{zp,i} &= \lim_{r \rightarrow \infty} \{ [t(\mathbf{r})\psi_i(\mathbf{r})] / \psi_i(\mathbf{r}) \} = -(1/2)\zeta_i^2 \\ &= \begin{cases} \epsilon_i & \text{for } s\text{-type atoms} \\ \epsilon_h & \text{otherwise} \end{cases} \end{aligned} \quad (4)$$

For s -type atoms, the correct Hartree-Fock orbital ψ_i must have the zero potential energy equivalent to its orbital energy. The orbitalwise criterion in this special case is identical to that for the Hartree orbital. For other cases, however, the zero potential energy should be equal to the orbital energy of the highest occupied orbital, independent of i ! This curious condition for the Hartree-Fock orbital originates from the nonvanishing contribution of exchange terms at $r \rightarrow \infty$. Indeed, we have for systems other than s -type atoms that

$$\kappa_i = \lim_{r \rightarrow \infty} \left\{ \left[\sum_{j(\neq i)} K_j(\mathbf{r}) \psi_j(\mathbf{r}) \right] / \psi_i(\mathbf{r}) \right\} = \epsilon_h - \epsilon_i, \quad (5)$$

where K_j is the exchange operator. At a point infinitely apart

from the nuclei, the exchange term κ_i locally has a constant contribution which is directly related to the orbital energy. Equation (4) is the consistency condition for the asymptotic behavior of the Hartree-Fock orbital and the associated orbital energy. Applications of this criterion will be presented in Sec. IV for a few atoms.

III. HARTREE-FOCK ZERO POTENTIAL ENERGY

The Hartree-Fock wave function is written as

$$\Psi(\{\mathbf{r}_i\}) = [(2N)!]^{-1/2} \sum_P (-1)^P P \times [\psi_1(1)\bar{\psi}_1(2)\cdots\psi_N(2N-1)\bar{\psi}_N(2N)], \quad (6)$$

where the symbols have the ordinary meaning. The zero potential energy for the wave function (6) is then

$$E_{zp} = \sum_i \left\{ \lim_{r_i \rightarrow \infty} [t(\mathbf{r}_i)\Psi(\{\mathbf{r}_i\})]/\Psi(\{\mathbf{r}_i\}) \right\}. \quad (7)$$

Using the asymptotic form of a spatial function [Eq. (3)], we find for a large r that

$$t(\mathbf{r}_i)\Psi(\{\mathbf{r}_i\}) = -(1/2)\zeta_i^2\Psi(\{\mathbf{r}_i\}) \quad (\text{large } r_i) \quad (8)$$

and therefore Eq. (7) is reduced to

$$E_{zp} = \begin{cases} 2 \sum_i \epsilon_i & \text{for } s\text{-type atoms} \\ 2N\epsilon_h & \text{otherwise} \end{cases} \quad (9)$$

Equation (9) shows that the Hartree-Fock zero potential energy is the sum of orbital energies of the occupied orbital or the orbital energy of the highest occupied orbital times the number of electrons. It constitutes the consistency condition for the Hartree-Fock wave function and the total electronic energy, deduced from the long-range asymptotic consideration. Evidently, E_{zp} does not agree with E_{av} which is given by

$$E_{av} = 2 \sum_i \epsilon_i - \sum_{i,j} (2J_{ij} - K_{ij}). \quad (10)$$

Moreover, it is quite unrealistic to assume that the true energy E , though it is unknown, is expressible in the form of Eq. (9). Thus, the Hartree-Fock wave function never satisfies the zero potential energy criterion [Eq. (2)], which accordingly implies that the Hartree-Fock approximation cannot correctly describe the long-range asymptotic behavior of many-electron wave functions. Though the Hartree-Fock approximation is a well-established independent-particle model and constitutes a beautiful theoretical framework, it may not be very suitable for problems such as scattering and long-range forces if accurate and careful descriptions of the wave function tail are needed. The same is true for the Hartree-Fock approximation.

IV. ILLUSTRATIONS

The majority of atomic and molecular Hartree-Fock calculations uses some basis functions and the (spatial) orbital is approximated by a linear combination of basis functions. Because of the required asymptotic form [Eq. (3)], basis functions other than the Slater and hydrogenic types have no chance to satisfy the zero potential energy criterion. For example, a Gaussian-type function $\chi(\mathbf{r}) = f(x,y,z)\exp(-\alpha r^2)$ gives $\epsilon_{zp} = -\infty$. For this reason, we here examine approximate Hartree-Fock wave functions obtained with Slater-type basis functions.

Table I summarizes the analysis of two types of ground state wave functions for He, Be, and Ne atoms. One is the unconstrained wave function due to Clementi and Roetti²⁰ and the other is the constrained one due to Weber *et al.*²¹ and Marron *et al.*,²² in which the orbitals are forced to satisfy $\zeta_i = (-2\epsilon_i)^{1/2}$ asymptotically. The latter authors considered $\zeta_i = (-2\epsilon_h)^{1/2}$ to be pathologic while $\zeta_i = (-2\epsilon_i)^{1/2}$ to be physically meaningful.

TABLE I. Zero potential energy criteria applied to the Hartree-Fock wave functions for the ground state of He, Be, and Ne atoms.

Atom	Orbital	Orbitalwise		Total		Zero potential energy criterion		
		$-\epsilon_i$	$-\epsilon_{zp,i}$	$-E_{av}$	$-E_{zp}$	Orbitalwise Eq. (4)	Hartree-Fock Eq. (9)	Exact Eq. (2)
Unconstrained wave functions ^a								
He	1s	0.917 95	1.004 14	2.861 68	2.008 29	No	No	No
Be	1s	4.732 67	0.302 80	14.573 02	1.211 19	No	No	No
	2s	0.309 27	0.302 80			No		
Ne	1s	32.772 48	1.924 41	128.547 05	14.023 24	No	No	No
	2s	1.930 43	1.924 41			No		
	2p	0.850 44	1.054 27					
Constrained wave functions ^b								
He	1s	0.917 96	0.917 96	2.861 68	1.835 92	Yes	Yes	No
Be	1s	4.732 79	4.732 80	14.573 01	10.084 16	Yes	Yes	No
	2s	0.309 29	0.309 28			Yes		
Ne	1s	32.773 34	32.773 34	128.546 19	74.513 68	No	No	No
	2s	1.930 50	1.930 57			No		
	2p	0.850 98	0.850 98			Yes		

^a Reference 20.

^b References 21 and 22.

The unconstrained wave functions reveal one general feature of basis-set-expansion wave functions: In the long-range asymptotic region, the basis function with the smallest exponent governs the behavior of an orbital unless its coefficient of the linear combination is exactly zero. Therefore, the different orbitals $\{\psi_i\}$ expressed by the same basis set have the identical $\epsilon_{zp,i}$. The situation is similar to the orbitalwise criterion [Eq. (4)], but orbitals with different symmetries (e.g., *s*- and *p*-type orbitals in Ne atom) can have different $\epsilon_{zp,i}$'s, since the exponents are not generally common to basis functions for different symmetries. However, the correct asymptotic behavior [Eq. (3)] suggests that the smallest exponent should be common to all basis sets for different symmetries and be equal to $(-2\epsilon_h)^{1/2}$, where ϵ_h is the energy of the highest occupied orbital. As a result, none of the unconstrained wave function in Table I fulfills the orbitalwise and Hartree-Fock zero potential energy criteria, though the 2*s* orbitals for Be and Ne show relatively good agreement of ϵ_i and $\epsilon_{zp,i}$.

On the other hand, the constrained wave functions for He and Be atoms satisfy both the orbitalwise and Hartree-Fock criteria. For Ne atom, however, only the 2*p* orbital satisfies the orbitalwise criterion, since $\zeta_i = (-2\epsilon_i)^{1/2}$ is used as the constraint.

We have also examined several Hartree-Fock wave functions for diatomic molecules calculated by Cade, Huo, and Wahl.²³⁻²⁵ The results are similar to those of the unconstrained atomic wave functions given in Table I and no detailed numerical data are presented here.

V. SUMMARY

We have shown that the exact zero potential energy criterion [Eq. (2)] cannot be satisfied by the Hartree-Fock wave function. This suggests the Hartree-Fock method may not be suitable for problems in which a very accurate description of the long-range asymptotic behavior is needed. The Hartree-Fock [Eq. (9)] and orbitalwise [Eq. (4)] criteria have been derived, which measure the consistency of the energy and the wave function tail at the level of the Hartree-Fock approximation. The present results are extended straightforwardly to the unrestricted Hartree-Fock wave function: The orbitalwise criterion applies separately to the

α and β spin orbitals and the Hartree-Fock criterion is expressed as the sum of two contributions, since each of the two Fock equations for the α and β spin orbitals has the same asymptotic structure with that for closed shells discussed here.

ACKNOWLEDGMENT

The author thanks Dr. K. Ohta for helpful discussion.

- ¹T. Koga, *J. Chem. Phys.* **83**, 6301 (1985).
- ²T. Koga and H. Sasaki, *Theor. Chim. Acta* **72**, 27 (1987).
- ³T. Koga, T. Uchiyama, and M. Uji-ie, *J. Chem. Phys.* **87**, 4025 (1987).
- ⁴J. H. Bartlett, *Phys. Rev.* **51**, 661 (1937).
- ⁵A. A. Frost, *J. Chem. Phys.* **10**, 240 (1942).
- ⁶A. A. Frost, R. E. Kellog, and E. C. Curtis, *Rev. Mod. Phys.* **32**, 313 (1960).
- ⁷B. H. Armstrong, *Bull. Am. Phys. Soc.* **9**, 401 (1964).
- ⁸R. J. Drachman, *Phys. Rev.* **136**, A641 (1964).
- ⁹A. J. Thakkar and V. H. Smith, Jr., *Phys. Rev. A* **18**, 841 (1978).
- ¹⁰T. Koga, K. Ohta, and A. J. Thakkar, *Phys. Rev. A* **37**, 1411 (1988).
- ¹¹N. C. Handy, M. T. Marron, and H. J. Silverstone, *Phys. Rev.* **180**, 45 (1969).
- ¹²G. S. Handler, D. W. Smith, and H. J. Silverstone, *J. Chem. Phys.* **73**, 3936 (1980).
- ¹³A. J. O'Connor, *Commun. Math. Phys.* **32**, 319 (1973).
- ¹⁴J. M. Combes and L. Thomas, *Commun. Math. Phys.* **34**, 251 (1973).
- ¹⁵E. H. Lieb and B. Simon, *Commun. Math. Phys.* **53**, 185 (1977).
- ¹⁶M. Hoffmann-Ostenhof and T. Hoffmann-Ostenhof, *Phys. Rev. A* **16**, 1782 (1977).
- ¹⁷R. Ahlrichs, M. Hoffmann-Ostenhof, and T. Hoffmann-Ostenhof, *J. Chem. Phys.* **68**, 1402 (1978).
- ¹⁸T. Hoffmann-Ostenhof, M. Hoffmann-Ostenhof, and R. Ahlrichs, *Phys. Rev. A* **18**, 328 (1978).
- ¹⁹S. H. Patil, *J. Chem. Phys.* **80**, 2689 (1984).
- ²⁰E. Clementi and C. Roetti, *At. Data Nucl. Data Tables* **14**, 177 (1974).
- ²¹T. A. Weber, N. C. Handy, and R. G. Parr, *J. Chem. Phys.* **52**, 1501 (1970).
- ²²M. T. Marron, N. C. Handy, R. G. Parr, and H. J. Silverstone, *Int. J. Quantum Chem.* **4**, 245 (1970).
- ²³P. E. Cade and W. Huo, *At. Data Nucl. Data Tables* **12**, 415 (1973).
- ²⁴P. E. Cade and A. C. Wahl, *At. Data Nucl. Data Tables* **13**, 339 (1974).
- ²⁵P. E. Cade and W. M. Huo, *At. Data Nucl. Data Tables* **15**, 1 (1975).