Envelopes of families of Legendre mappings in the unit tangent bundle over the Euclidean space

著者	TAKAHASHI Masatomo
journal or publication title	Journal of Mathematical Analysis and Applications
volume	473
number	1
page range	408-420
year	2019-05-01
URL	http://hdl.handle.net/10258/00009919

doi: info:doi/10.1016/j.jmaa.2018.12.057

Envelopes of families of Legendre mappings in the unit tangent bundle over the Euclidean space

Masatomo Takahashi

Dedicated to Professor Takashi Nishimura on the occasion of his 60th birthday

November 7, 2018

Abstract

For families of hypersurfaces with singular points, a classical definition of an envelope is vague. In order to define an envelope for a family of hypersurfaces with singular points, we consider r -parameter families of frontals and of Legendre mappings in the unit tangent bundle over the Euclidean space. We define an envelope for the r -parameter family of Legendre mappings. Then the envelope is also a frontal. We investigate properties of the envelopes. As an application, we give a condition that the projection of a singular solution of a first order partial differential equation is an envelope.

1 Introduction

Envelopes are classical object in the differential geometry. There are a lot of applications of envelopes to differential geometry, differential equations and physics, for instance [4, 5, 7, 8, 9, 12, 15, 16, 18, 21, 23]. An envelope of a family of surfaces is a surface that is "tangent" to each member of the family at some point. If the surfaces are regular, then the tangent is well-defined. However, a definition of an envelope is vague for singular surfaces (surfaces with singular points). In [22], a definition and properties of an envelope for a one-parameter family of Legendre curves in the unit tangent bundle over $\mathbb{R}^2$ were given. In this paper, we clarify a definition of an envelope for a family of singular surfaces. As singular surfaces, we consider frontals and Legendre mappings in the unit tangent bundle over $\mathbb{R}^{n+1}$. The basic results on the singularity theory see [1, 2, 4, 13, 14, 17].

We consider r -parameter families of Legendre mappings and define an envelope in §3. Then the envelope of an r -parameter family of Legendre mappings is also a frontal. We give a necessary and sufficient condition that the r -parameter family of Legendre mappings has an envelope, see Theorem 3.6 as the envelope theorem. Moreover, we give relations between envelopes of a classical definition and of a family of Legendre mappings. As an application,

2010 Mathematics Subject classification: 58K05, 53A07, 57R45

Key Words and Phrases. envelope, family of Legendre mappings, frontal, singular solution.

we give a condition that the projection of a singular solution of a first order partial differential equation is an envelope by using the envelope theorem in §4.

All maps and manifolds considered here are differentiable of class C^∞ .

Acknowledgement. The author would like to thank Professor Takashi Nishimura for his constant encouragements. The author would also like to thank Professor Kentaro Saji for valuable discussions and the referee for useful comments that improved the original manuscript. This work was partially supported by JSPS KAKENHI Grant Number JP 17K05238.

2 Preliminaries

Let $\mathbb{R}^{n+1}$ be the $(n+1)$ -dimensional Euclidean space with the inner product $x \cdot y = x_1y_1 + \cdots + x_{n+1}y_{n+1}$, where $x = (x_1, \dots, x_{n+1})$, $y = (y_1, \dots, y_{n+1}) \in \mathbb{R}^{n+1}$. The norm of $x \in \mathbb{R}^{n+1}$ is given by $|x| = \sqrt{x \cdot x}$.

Let $F : W \times \Lambda \rightarrow \mathbb{R}$ be an r -parameter family of smooth functions, where W and Λ are domains in $\mathbb{R}^{n+1}$ and in $\mathbb{R}^r$, respectively. Then one of the classical definition of an envelope E_I is as follows, see for instance [3, 4, 11]:

Definition 2.1 The *envelope* of the family F is the discriminant set of F , that is, the set of points given by

$$E_I = \{x \in \mathbb{R}^{n+1} \mid \text{for some } \lambda \in \Lambda, F(x, \lambda) = F_{\lambda_j}(x, \lambda) = 0, j = 1, \dots, r\}.$$

If $F(x, \lambda) = F_{\lambda_j}(x, \lambda) = 0$, $j = 1, \dots, r$, we say that $x \in E_I$ with respect to $\lambda = (\lambda_1, \dots, \lambda_r)$. Here $F_{\lambda_j}(x, \lambda) = (\partial F / \partial \lambda_j)(x, \lambda)$.

Example 2.2 Let $F : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{R}$, $F(x, y, z, \lambda) = (x - \lambda)^3 - y^2$. Then $F = 0$ is the image of the cuspidal edge for each fixed $\lambda \in \mathbb{R}$, see Figure 1 and Example 3.8. The definition and properties of cuspidal edges see [10, 20]. Since $F_\lambda(x, y, z, \lambda) = -3(x - \lambda)^2$, the envelope of the family F is given by $E_I = \{(\lambda, 0, z)\} = xz$ -plane.

Example 2.3 Let $F : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{R}$, $F(x, y, z, \lambda) = x^3 - (y - \lambda)^2$. Then $F = 0$ is the image of the cuspidal edge for each fixed $\lambda \in \mathbb{R}$, see Figure 2 and Example 3.9. Since $F_\lambda(x, y, z, \lambda) = 2(y - \lambda)$, the envelope of the family F is given by $E_I = \{(0, \lambda, z)\} = yz$ -plane.

Figure 1.

Figure 2.

However, in the sense of the limit tangent plane of the cuspidal edge, yz -plane is not tangent to the cuspidal edge. Therefore, we would like to distinguish as envelopes, see Examples 3.8 and 3.9.

Let $U \subset \mathbb{R}^n$ be a domain in $\mathbb{R}^n$. We say that $(f, \nu) : U \rightarrow \mathbb{R}^{n+1} \times S^n$ is a *Legendre mapping* if $(f, \nu)^*\theta = 0$, where θ is a canonical contact form on the unit tangent bundle $T_1\mathbb{R}^{n+1} = \mathbb{R}^{n+1} \times S^n$ over $\mathbb{R}^{n+1}$ (cf. [1, 2]). Moreover, $f : U \rightarrow \mathbb{R}^{n+1}$ is a *frontal* (respectively, a *front*) if there exists a smooth mapping $\nu : U \rightarrow S^n$ such that (f, ν) is a Legendre mapping (respectively, a Legendre immersion). The condition $(f, \nu)^*\theta = 0$ is equivalent to $df(u) \cdot \nu(u) = 0$ for all $u \in U$. If we denote $f(u) = (f_1(u), \dots, f_{n+1}(u))$, $\nu(u) = (\nu_1(u), \dots, \nu_{n+1}(u))$ and $u = (u_1, \dots, u_n)$, then the condition $df(u) \cdot \nu(u) = 0$ for all $u \in U$ is equivalent to

$$f_{u_i}(u) \cdot \nu(u) = f_{1u_i}(u)\nu_1(u) + \dots + f_{n+1u_i}(u)\nu_{n+1}(u) = 0,$$

for all $u \in U$ and $i = 1, \dots, n$.

The *parallel* of a Legendre mapping $(f, \nu) : U \rightarrow \mathbb{R}^{n+1} \times S^n$ is defined by $f^k : U \rightarrow \mathbb{R}^{n+1}$, $f^k(u) = f(u) + k\nu(u)$, where $k \in \mathbb{R}$. Then it is easy to see that $(f^k, \nu) : U \rightarrow \mathbb{R}^{n+1} \times S^n$ is also a Legendre mapping for each fixed $k \in \mathbb{R}$.

3 Envelopes of families of Legendre mappings

We say that $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ is an *r -parameter family of Legendre mapping* if $(f(\cdot, \lambda), \nu(\cdot, \lambda)) : U \rightarrow \mathbb{R}^{n+1} \times S^n$ is a Legendre mapping for each $\lambda \in \Lambda \subset \mathbb{R}^r$.

Let $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ be an r -parameter family of Legendre mappings. Let $V \subset \mathbb{R}^n$ be an open subset and $e : V \rightarrow U \times \Lambda$, $e(p) = (u(p), \lambda(p))$ be a smooth mapping. We denote $E = f \circ e : V \rightarrow \mathbb{R}^{n+1}$.

Definition 3.1 We call E an *envelope* (and e a *pre-envelope*) for the r -parameter family of Legendre mappings (f, ν) , when the following conditions are satisfied.

- (i) The set of regular points of $\lambda : V^n \rightarrow \Lambda^r$ is dense in V . (The Variability Condition.)
- (ii) For all $p \in V$ and $i = 1, \dots, n$, $E_{p_i}(p) \cdot \nu(e(p)) = 0$. (The Tangency Condition.)

The definition of the envelope is a generalisation of the definition of the envelope of a one-parameter family of Legendre curves in [22]. By definition, we have the following.

Proposition 3.2 Let $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ be an r -parameter family of Legendre mappings. Suppose that $e : V \rightarrow U \times \Lambda$ is a pre-envelope and $E = f \circ e : V \rightarrow \mathbb{R}^{n+1}$ is an envelope of (f, ν) . Then E is a frontal. More precisely, $(E, \nu \circ e) : V \rightarrow \mathbb{R}^{n+1} \times S^n$ is a Legendre mapping.

Proof. Since the tangency condition, we have $E_{p_i}(p) \cdot \nu(e(p)) = 0$ for all $p \in V$. It follows that $dE(p) \cdot (\nu \circ e)(p) = 0$ for all $p \in V$. That is, $(E, \nu \circ e) : V \rightarrow \mathbb{R}^{n+1} \times S^n$ is a Legendre mapping. $\square$

Proposition 3.3 Let $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ be an r -parameter family of Legendre mappings. Suppose that $e : V \rightarrow U \times \Lambda$ is a pre-envelope and $E = f \circ e$ is an envelope of (f, ν) . Then we have the following.

(1) $e : V \rightarrow U \times \Lambda$ is also a pre-envelope of $(f, -\nu)$ and $E = f \circ e$ is also an envelope of $(f, -\nu)$.

(2) $e : V \rightarrow U \times \Lambda$ is also a pre-envelope of $(-f, \nu)$ and $-E = -f \circ e$ is an envelope of $(-f, \nu)$.

Proof. (1) By definition, $(f, -\nu)$ is also an r -parameter family of Legendre mappings. Since e is a pre-envelope of (f, ν) , $E_{p_i}(p) \cdot (-\nu(e(p))) = -E_{p_i}(p) \cdot \nu(e(p)) = 0$ for all $p \in V$. Hence, e is also a pre-envelope and $E = f \circ e$ is also an envelope of $(f, -\nu)$.

(2) By similarly, we have the result. $\square$

Remark 3.4 By Proposition 3.3 (1), we may define an envelope for an r -parameter family of Legendre mapping in $PT^*\mathbb{R}^{n+1}$.

Remark 3.5 As the same definition, we can define an envelope of a family of Legendre mappings in the unit tangent bundle over a smooth manifold. Especially, we can define envelopes not only of families of Legendre mappings in the unit spherical bundle (cf. [19]), but also of families of frontals in the hyperbolic or de-Sitter space (cf. [6]).

We give a necessary and sufficient condition that the r -parameter family of Legendre mappings has an envelope. We call this result *the envelope theorem* (cf. [11, 22]).

Theorem 3.6 (The Envelope Theorem) *Let $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ be an r -parameter family of Legendre mappings, and let $e : V \rightarrow U \times \Lambda$ be a smooth mapping satisfying the variability condition. Suppose that $n \geq r$. Then e is a pre-envelope of (f, ν) (and E is an envelope) if and only if $f_{\lambda_j}(e(p)) \cdot \nu(e(p)) = 0$ for all $p \in V$ and $j = 1, \dots, r$.*

Proof. Suppose that e is a pre-envelope of (f, ν) . We denote $f = (f_1, \dots, f_{n+1}), \nu = (\nu_1, \dots, \nu_{n+1})$. By a direct calculation,

$$\begin{aligned} E_{p_i}(p) &= \frac{\partial}{\partial p_i}(f \circ e(p)) \\ &= \left(\sum_{k=1}^n f_{1u_k}(e(p))u_{kp_i}(p) + \sum_{j=1}^r f_{1\lambda_j}(e(p))\lambda_{jp_i}(p), \dots, \right. \\ &\quad \left. \sum_{k=1}^n f_{n+1u_k}(e(p))u_{kp_i}(p) + \sum_{j=1}^r f_{n+1\lambda_j}(e(p))\lambda_{jp_i}(p) \right). \end{aligned}$$

Since $E_{p_i}(p) \cdot \nu(e(p)) = 0$ for all $p \in V$ and $i = 1, \dots, n$, and (f, ν) is an r -parameter family of Legendre mappings, we have

$$(f_{\lambda_1}(e(p)) \cdot \nu(e(p)))\lambda_{1p_i}(p) + \dots + (f_{\lambda_r}(e(p)) \cdot \nu(e(p)))\lambda_{rp_i}(p) = 0,$$

for all $p \in V$ and $i = 1, \dots, n$. It follows that

$$\begin{pmatrix} \lambda_{1p_1}(p) & \cdots & \lambda_{rp_1}(p) \\ \vdots & \cdots & \vdots \\ \lambda_{1p_n}(p) & \cdots & \lambda_{rp_n}(p) \end{pmatrix} \begin{pmatrix} f_{\lambda_1}(e(p)) \cdot \nu(e(p)) \\ \vdots \\ f_{\lambda_r}(e(p)) \cdot \nu(e(p)) \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}.$$

By the assumption $n \geq r$ and the variability condition, we have $f_{\lambda_j}(e(p)) \cdot \nu(e(p)) = 0$ for all $p \in V$ and $j = 1, \dots, r$.

Conversely, suppose that $f_{\lambda_j}(e(p)) \cdot \nu(e(p)) = 0$ for all $p \in V$ and $j = 1, \dots, r$. By a direct calculation, we have

$$\begin{aligned}
E_{p_i}(p) \cdot \nu(e(p)) &= \left(\sum_{k=1}^n f_{1u_k}(e(p)) u_{kp_i}(p) + \sum_{j=1}^r f_{1\lambda_j}(e(p)) \lambda_{jp_i}(p) \right) \cdot \nu_1(e(p)) \\
&\quad + \dots + \left(\sum_{k=1}^n f_{n+1u_k}(e(p)) u_{kp_i}(p) + \sum_{j=1}^r f_{n+1\lambda_j}(e(p)) \lambda_{jp_i}(p) \right) \cdot \nu_{n+1}(e(p)) \\
&= \sum_{k=1}^n u_{kp_i}(p) f_{u_k}(e(p)) \cdot \nu(e(p)) + \sum_{j=1}^r \lambda_{jp_i}(p) f_{\lambda_j}(e(p)) \cdot \nu(e(p)) \\
&= 0
\end{aligned}$$

for all $p \in V$ and $i = 1, \dots, n$. It follows that e is a pre-envelope of (f, ν) . $\square$

Remark 3.7 In Theorem 3.6, the assumption $n \geq r$ does not need to prove the converse.

Example 3.8 Let $(f, \nu) : \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}^3 \times S^2$ be

$$f(u, v, \lambda) = (u^2 + \lambda, u^3, v), \nu(u, v, \lambda) = \frac{1}{\sqrt{9u^2 + 4}}(3u, -2, 0).$$

Then (f, ν) is a one-parameter family of Legendre mappings (immersions) and f is the cuspidal edge for each fixed $\lambda \in \mathbb{R}$. Since $f_{\lambda}(u, v, \lambda) \cdot \nu(u, v, \lambda) = 3u/\sqrt{9u^2 + 4}$, if we take $e : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \times \mathbb{R}$, $e(p, q) = (0, p, q)$, then the variability condition holds and $f_{\lambda}(e(p, q)) \cdot \nu(e(p, q)) = 0$ for all $(p, q) \in \mathbb{R}^2$. By Theorem 3.6, e is a pre-envelope and $E(p, q) = f \circ e(p, q) = (q, 0, p)$ is an envelope. Hence xz -plane is an envelope of (f, ν) , see Example 2.2.

Example 3.9 Let $(f, \nu) : \mathbb{R}^2 \times \mathbb{R} \rightarrow \mathbb{R}^3 \times S^2$ be

$$f(u, v, \lambda) = (u^2, u^3 + \lambda, v), \nu(u, v, \lambda) = \frac{1}{\sqrt{9u^2 + 4}}(3u, -2, 0).$$

Then (f, ν) is a one-parameter family of Legendre mappings (immersions) and f is the cuspidal edge for each fixed $\lambda \in \mathbb{R}$. Since $f_{\lambda}(u, v, \lambda) \cdot \nu(u, v, \lambda) = -2/\sqrt{9u^2 + 4} \neq 0$, (f, ν) does not have an envelope by Theorem 3.6. Hence yz -plane is not an envelope of (f, ν) , see Example 2.3.

Definition 3.10 We say that a map $\Phi : \tilde{U} \times \tilde{\Lambda} \rightarrow U \times \Lambda$ is an r -parameter family of parameter change if Φ is a diffeomorphism and given by the form $\Phi(q, k) = (\phi(q, k), \varphi(k))$.

Proposition 3.11 Let $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ be an r -parameter family of Legendre mappings. Suppose that $n \geq r$, $e : V \rightarrow U \times \Lambda$ is a pre-envelope, $E = f \circ e$ is an envelope and $\Phi : \tilde{U} \times \tilde{\Lambda} \rightarrow U \times \Lambda$ is an r -parameter family of parameter change. Then $(\tilde{f}, \tilde{\nu}) = (f \circ \Phi, \nu \circ \Phi) : \tilde{U} \times \tilde{\Lambda} \rightarrow \mathbb{R}^{n+1} \times S^n$ is also an r -parameter family of Legendre mappings. Moreover, $\Phi^{-1} \circ e : V \rightarrow \tilde{U} \times \tilde{\Lambda}$ is a pre-envelope and E is also an envelope of $(\tilde{f}, \tilde{\nu})$.

Proof. By the chain rule, we have $d(f \circ \Phi) \cdot \nu \circ \Phi = df(\Phi)d\Phi \cdot \nu(\Phi) = 0$ for fixed $k \in \tilde{\Lambda}$. Therefore, $(\tilde{f}, \tilde{\nu})$ is also an r -parameter family of Legendre mappings. Since the form of Φ ,

there exists a smooth map $\psi : U \times \Lambda \rightarrow \tilde{U}$ such that $\Phi^{-1}(u, \lambda) = (\psi(u, \lambda), \varphi^{-1}(\lambda))$. It follows that $\Phi^{-1} \circ e$ satisfies the variability condition. By a direct calculation, we have

$$\begin{aligned} \tilde{f}_{k_j}(q, k) &= \frac{\partial}{\partial k_j} f \circ \Phi(q, k) \\ &= \left(\sum_{i=1}^n f_{1u_i}(\Phi(q, k)) \phi_{ik_j}(q, k) + \sum_{\ell=1}^r f_{1\lambda_\ell}(\Phi(q, k)) \varphi_{\ell k_j}(k), \dots, \right. \\ &\quad \left. \sum_{i=1}^n f_{n+1u_i}(\Phi(q, k)) \phi_{ik_j}(q, k) + \sum_{\ell=1}^r f_{n+1\lambda_\ell}(\Phi(q, k)) \varphi_{\ell k_j}(k) \right) \end{aligned}$$

for all $(q, k) \in \tilde{U} \times \tilde{\Lambda}$ and $j = 1, \dots, r$. Then

$$\tilde{f}_{k_j}(\Phi^{-1} \circ e(p)) \cdot \tilde{\nu}(\Phi^{-1} \circ e(p)) = \varphi_{1k_j}(\lambda(p)) f_{\lambda_1}(e(p)) \cdot \nu(e(p)) + \dots + \varphi_{rk_j}(\lambda(p)) f_{\lambda_r}(e(p)) \cdot \nu(e(p)) = 0$$

for all $p \in V$ and $j = 1, \dots, r$. It follows that $\Phi^{-1} \circ e$ is a pre-envelope of $(\tilde{f}, \tilde{\nu})$, and hence $\tilde{f} \circ \Phi^{-1} \circ e = f \circ \Phi \circ \Phi^{-1} \circ e = f \circ e = E$ is also an envelope of $(\tilde{f}, \tilde{\nu})$. $\square$

Let $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ be an r -parameter family of Legendre mappings. We define the parallel of the r -parameter family of Legendre mappings by $f^k : U \times \Lambda \rightarrow \mathbb{R}^{n+1}$, $f^k(u, \lambda) = f(u, \lambda) + k\nu(u, \lambda)$, where $k \in \mathbb{R}$. It is easy to see that (f^k, ν) is also an r -parameter family of Legendre mappings for each fixed $k \in \mathbb{R}$.

Proposition 3.12 *Suppose that $e : V \rightarrow U \times \Lambda$ is a pre-envelope of (f, ν) (and E is an envelope) and $n \geq r$. Then the envelope of the parallel of the r -parameter family of Legendre mappings is given by the parallel of the envelope.*

Proof. Since ν is a unit vector, $\nu_{\lambda_j}(u, \lambda) \cdot \nu(u, \lambda) = 0$. Therefore, $f_{\lambda_j}^k(u, \lambda) \cdot \nu(u, \lambda) = (f_{\lambda_j}(u, \lambda) + k\nu_{\lambda_j}(u, \lambda)) \cdot \nu(u, \lambda) = f_{\lambda_j}(u, \lambda) \cdot \nu(u, \lambda)$. If e is a pre-envelope of (f, ν) , then $f_{\lambda_j}^k(e(p)) \cdot \nu(e(p)) = f_{\lambda_j}(e(p)) \cdot \nu(e(p)) = 0$ for all $p \in V$ and $j = 1, \dots, r$. It follows that e is also a pre-envelope of (f^k, ν) by Theorem 3.6. By definition, the envelope of the parallel of the r -parameter family of Legendre mappings is given by $E^k = f^k \circ e = f \circ e + k\nu \circ e = E + k\nu \circ e$. It follows that E^k is the parallel of the Legendre mapping $(E, \nu \circ e)$. $\square$

We give a relation between the envelope E_I of the classical definition by using an implicit function (Definition 2.1) and the envelope E of an r -parameter family of Legendre mappings (Definition 3.1).

Proposition 3.13 *Let $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ be an r -parameter family of Legendre mappings, and let $F(x, \lambda) = 0$ be an implicit function of the r -parameter family of frontals, that is, assume $F(f(u, \lambda), \lambda) = 0$ and $(F_{x_1}, \dots, F_{x_{n+1}})(f(u, \lambda), \lambda)$ is parallel to $\nu(u, \lambda)$ for all $(u, \lambda) \in U \times \Lambda$. Suppose that $n \geq r$. If $e : V \rightarrow U \times \Lambda$ is a pre-envelope and $E : V \rightarrow \mathbb{R}^{n+1}$ is an envelope of (f, ν) , then $E(V) \subset E_I$.*

Proof. By differentiating $F(f(u, \lambda), \lambda) = 0$ with respect to λ_j , we have

$$F_{x_1}(f(u, \lambda), \lambda) f_{1\lambda_j}(u, \lambda) + \dots + F_{x_{n+1}}(f(u, \lambda), \lambda) f_{n+1\lambda_j}(u, \lambda) + F_{\lambda_j}(f(u, \lambda), \lambda) = 0,$$

where $j = 1, \dots, r$. By the assumption, there exists a smooth function $a : U \times \Lambda \rightarrow \mathbb{R}$ such that $(F_{x_1}, \dots, F_{x_{n+1}})(f(u, \lambda), \lambda) = a(u, \lambda)(\nu_1, \dots, \nu_{n+1})(u, \lambda)$ for all $(u, \lambda) \in U \times \Lambda$. By

Theorem 3.6, we have $f_{\lambda_j}(e(p)) \cdot \nu(e(p)) = 0$ for all $p \in V$ and $j = 1, \dots, r$. It follows that $F_{\lambda_j}(f(e(p)), \lambda(p)) = 0$ for all $p \in V$ and $j = 1, \dots, r$. Therefore $E(p) \in E_I$ with respect to $\lambda(p)$ for all $p \in V$. $\square$

Proposition 3.14 *Let $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ be an r -parameter family of Legendre mappings, and let $e : V \rightarrow U \times \Lambda$ be a smooth map satisfying the variability condition. If $\text{rank}(f_{u_1}, \dots, f_{u_n})(e(p)) = n$ and the trace of e lies in the singular set of f , then e is a pre-envelope of (f, ν) (and E is an envelope).*

Proof. We denote the set of singular points (singular set) of f by $\Sigma(f)$. Since $e(p) \in \Sigma(f)$, we have the condition

$$\text{rank} \begin{pmatrix} f_{1u_1} & \cdots & f_{1u_n} & f_{1\lambda_1} & \cdots & f_{1\lambda_r} \\ \vdots & & \vdots & \vdots & & \vdots \\ f_{n+1u_1} & \cdots & f_{n+1u_n} & f_{n+1\lambda_1} & \cdots & f_{n+1\lambda_r} \end{pmatrix} (e(p)) < n + 1.$$

By the assumption $\text{rank}(f_{u_1}, \dots, f_{u_n})(e(p)) = n$, there exist smooth functions $a_{ij} : V \rightarrow \mathbb{R}$, $i = 1, \dots, n, j = 1, \dots, r$ such that $f_{\lambda_j}(e(p)) = a_{1j}(p)f_{u_1}(e(p)) + \cdots + a_{nj}(p)f_{u_n}(e(p))$. It follows that $f_{\lambda_j}(e(p)) \cdot \nu(e(p)) = 0$ for all $p \in V$ and $j = 1, \dots, r$. Hence e is a pre-envelope of (f, ν) . $\square$

Proposition 3.15 *Let $(f, \nu) : U \times \Lambda \rightarrow \mathbb{R}^{n+1} \times S^n$ be an r -parameter family of Legendre mappings, and let $F(x, \lambda) = 0$ be an implicit function of the r -parameter family of frontals, that is, assume $F(f(x, \lambda), \lambda) = 0$ and $(F_{x_1}, \dots, F_{x_{n+1}})(f(u, \lambda), \lambda)$ is parallel to $\nu(u, \lambda)$ for all $(u, \lambda) \in U \times \Lambda$. Suppose that $e : V \rightarrow U \times \Lambda, e(p) = (u(p), \lambda(p))$ is a smooth mapping satisfying the variability condition. If $E(p) = f \circ e(p) \in E_I$ with respect to $\lambda(p)$, $\text{rank}(f_{u_1}, \dots, f_{u_n})(e(p)) = n$ and*

$$(F_{x_1}, \dots, F_{x_{n+1}})(f(e(p)), \lambda(p)) \neq (0, \dots, 0)$$

for all $p \in V$, then e is a pre-envelope of (f, ν) (and E is an envelope).

Proof. By differentiating $F(f(u, \lambda), \lambda) = 0$ with respect to u_i and λ_j , we have

$$\begin{aligned} F_{x_1}(f(u, \lambda), \lambda)f_{1u_i}(u, \lambda) + \cdots + F_{x_{n+1}}(f(u, \lambda), \lambda)f_{n+1u_i}(u, \lambda) &= 0, \\ F_{x_1}(f(u, \lambda), \lambda)f_{1\lambda_j}(u, \lambda) + \cdots + F_{x_{n+1}}(f(u, \lambda), \lambda)f_{n+1\lambda_j}(u, \lambda) + F_{\lambda_j}(f(u, \lambda), \lambda) &= 0, \end{aligned}$$

where $i = 1, \dots, n, j = 1, \dots, r$. Since $E(p) \in E_I$ with respect to $\lambda(p)$, we have $F_{\lambda_j}(f(e(p)), \lambda(p)) = 0$ for all $p \in V$ and $j = 1, \dots, r$. It follows that

$$\begin{pmatrix} f_{1u_1}(e(p)) & \cdots & f_{n+1u_1}(e(p)) \\ \vdots & & \vdots \\ f_{1u_n}(e(p)) & \cdots & f_{n+1u_n}(e(p)) \\ f_{1\lambda_1}(e(p)) & \cdots & f_{n+1\lambda_1}(e(p)) \\ \vdots & & \vdots \\ f_{1\lambda_r}(e(p)) & \cdots & f_{n+1\lambda_r}(e(p)) \end{pmatrix} \begin{pmatrix} F_{x_1}(f(e(p)), \lambda(p)) \\ \vdots \\ F_{x_{n+1}}(f(e(p)), \lambda(p)) \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}.$$

If the $\text{rank} df(e(p)) = n+1$, then $(F_{x_1}, \dots, F_{x_{n+1}})(f(e(p)), \lambda(p)) = (0, \dots, 0)$. By the assumption $(F_{x_1}, \dots, F_{x_{n+1}})(f(e(p)), \lambda(p)) \neq (0, \dots, 0)$, we have $\text{rank} df(e(p)) < n + 1$. It follows that $e(p) \in \Sigma(f)$. By Proposition 3.14, e is a pre-envelope of (f, ν) . $\square$

4 Singular solutions of first order partial differential equations

As an application of the theory of envelopes of families of Legendre mappings, we give a condition that the projection of a singular solution of a first order partial differential equation is an envelope.

We quickly review the theory of singular solutions and of Clairaut type of first order partial differential equations, in detail see [15, 16].

An equation is a submersion germ $F : (J^1(\mathbb{R}^n, \mathbb{R}), z_0) \rightarrow (\mathbb{R}, 0)$ on the 1-jet space of functions of n -variables, where $z_0 = (x_0, y_0, p_0)$. Let θ be a canonical contact 1-form on $J^1(\mathbb{R}^n, \mathbb{R})$ which is given by $\theta = dy - \sum_{i=1}^n p_i dx_i$, where $(x, y, p) = (x_1, \dots, x_n, y, p_1, \dots, p_n)$ is the canonical coordinate on $J^1(\mathbb{R}^n, \mathbb{R})$. We define a *geometric solution* of $F = 0$ to be an immersion germ $i : (L, q_0) \rightarrow (F^{-1}(0), z_0)$ of an n -dimensional manifold such that $i^*\theta = 0$, that is, a Legendre submanifold which is contained in $F^{-1}(0)$. We say that z_0 is a *contact singular point* if $\theta(T_{z_0}F^{-1}(0)) = 0$. It is easy to see that z_0 is a contact singular point if and only if $F = F_{p_i} = F_{x_i} + p_i F_y = 0$ for $i = 1, \dots, n$ at z_0 . We also say that z_0 is a π -singular point if $F = F_{p_i} = 0$ for $i = 1, \dots, n$ at z_0 . We denote the set of contact singular points by $\Sigma_c(F)$, the set of π -singular points by $\Sigma_\pi(F)$ and $\pi(\Sigma_\pi(F)) = D_F$, where $\pi : J^n(\mathbb{R}^n, \mathbb{R}) \rightarrow \mathbb{R}^{n+1}$ is the canonical projection $\pi(x, y, p) = (x, y)$. We call the set D_F a *discriminant set* of the equation $F = 0$.

An equation $F = 0$ is said to be *completely integrable* at z_0 if there exists a foliation by geometric solution on $F^{-1}(0)$ around z_0 , that is, there exists an immersion germ $\Gamma : (\mathbb{R}^n \times \mathbb{R}^n, (u_0, c_0)) \rightarrow (F^{-1}(0), z_0)$ such that $\Gamma(\cdot, c)$ is a geometric solution of $F = 0$ for each $c \in (\mathbb{R}^n, c_0)$. In this case, such a foliation is called a *complete solution* of $F = 0$ at z_0 . We say that an n -parameter family of function germs $f : (\mathbb{R}^n \times \mathbb{R}^n, (x_0, c_0)) \rightarrow (\mathbb{R}, y_0)$ is a *classical complete solution* of $F = 0$ at z_0 if a complete solution is a form of $j^1 f : (\mathbb{R}^n \times \mathbb{R}^n, (x_0, c_0)) \rightarrow (F^{-1}(0), z_0)$, that is, $F(x, f(x, c), f_x(x, c)) = 0$ and $j^1 f(x, c) = (x, f(x, c), f_x(x, c))$ is an immersion germ. An equation $F = 0$ is said to be *classical completely integrable* at z_0 if there exists a classical complete solution of $F = 0$ at z_0 .

A geometric solution $i : (L, q_0) \rightarrow (F^{-1}(0), z_0)$ of $F = 0$ is called a *singular solution* of $F = 0$ at z_0 if for any representative $\tilde{i} : U \rightarrow F^{-1}(0)$ of i and any open subset $V \subset U$, $\tilde{i}(V)$ is not contained in a leaf of any complete solutions of $F = 0$.

An equation $F = 0$ is called of *Clairaut type* at z_0 if there exist smooth function germs $B_{ji}, A_i : (J^1(\mathbb{R}^n, \mathbb{R}), z_0) \rightarrow \mathbb{R}$ for $i, j = 1, \dots, n$ such that

$$F_{x_i} + p_i F_y = \sum_{j=1}^n B_{ji} F_{p_j} + A_i F, \quad B_{ji} = B_{ij}$$

and

$$\frac{\partial B_{jk}}{\partial x_i} + p_i \frac{\partial B_{jk}}{\partial y} + \sum_{\ell=1}^n B_{\ell i} \frac{\partial B_{jk}}{\partial p_\ell} = \frac{\partial B_{ji}}{\partial x_k} + p_k \frac{\partial B_{ji}}{\partial y} + \sum_{\ell=1}^n B_{\ell k} \frac{\partial B_{ji}}{\partial p_\ell}$$

at any $(x, y, p) \in (F^{-1}(0), z_0)$ for $i, j, k = 1, \dots, n$. Then we have the following result.

Theorem 4.1 ([15, 16]) *Let $F : (J^1(\mathbb{R}^n, \mathbb{R}), z_0) \rightarrow (\mathbb{R}, 0)$ be a first order partial differential equation germs.*

(1) $F = 0$ is completely integrable at z_0 if and only if $\Sigma_c(F) = \emptyset$ or $\Sigma_c(F)$ is an n -dimensional submanifold around z_0 . Moreover, if $\Sigma_c(F) \neq \emptyset$, then $\Sigma_c(F)$ is a singular solution of $F = 0$ at z_0 .

(2) $F = 0$ is smooth completely integrable at z_0 if and only if $F = 0$ is of Clairaut type at z_0 . In this case, if $\Sigma_\pi(F) \neq \emptyset$, then $\Sigma_\pi(F)$ is a singular solution of $F = 0$ at z_0 and the discriminant set D_F is the envelope of the family of graphs of the smooth complete solution.

By using the envelope theorem (Theorem 3.6), we have the following result.

Theorem 4.2 Let $F : (J^1(\mathbb{R}^n, \mathbb{R}), z_0) \rightarrow (\mathbb{R}, 0)$ be a first order partial differential equation germs and not of Clairaut type at z_0 . Suppose that $\Gamma = (x, y, p) : (\mathbb{R}^n \times \mathbb{R}^n, (u_0, c_0)) \rightarrow (F^{-1}(0), z_0)$ is a complete solution of $F = 0$ at z_0 , $\Sigma_c(F) = \Sigma_\pi(F) \neq \emptyset$ and $e : (\mathbb{R}^n, \tilde{q}_0) \rightarrow (\mathbb{R}^n \times \mathbb{R}^n, (u_0, c_0))$ is a smooth mapping satisfying the variability condition. Then e is a pre-envelope and $E = \pi \circ \Gamma \circ e$ is an envelope of $(\pi \circ \Gamma, \nu)$ if and only if $E(q) \in \pi(\Sigma_c(F))$ for all $q \in (\mathbb{R}^n, \tilde{q}_0)$, where $\nu(u, c) = (-p(u, c), 1)/\sqrt{1 + |p(u, c)|^2}$.

Proof. By the assumption and Theorem 4.1 (1), $\Sigma_c(F) = \Sigma_\pi(F)$ is an n -dimensional manifold around z_0 and a singular solution of $F = 0$ at z_0 . Since $z_0 \in \Sigma_c(F)$ and $F = 0$ is submersion, we may consider $F(x, y, p) = -y + g(x, p)$, where g is a smooth function, $x = (x_1, \dots, x_n)$ and $p = (p_1, \dots, p_n)$. We denote the complete solution of $F = 0$ at z_0 by

$$\Gamma(u, c) = (x(u, c), y(u, c), p(u, c)) = (x_1(u, c), \dots, x_n(u, c), y(u, c), p_1(u, c), \dots, p_n(u, c)),$$

where $u = (u_1, \dots, u_n)$, $c = (c_1, \dots, c_n)$. Since $y(u, c) = g(x(u, c), p(u, c))$ and $\Gamma^*\theta = 0$, we have $y_{u_i}(u, c) = p_1(u, c)x_{1u_i}(u, c) + \dots + p_n(u, c)x_{nu_i}(u, c)$ for $i = 1, \dots, n$. Since $F = 0$ is not of Clairaut type at z_0 , we have

$$\text{rank} \begin{pmatrix} x_{1u_1} & \cdots & x_{nu_1} \\ \vdots & & \vdots \\ x_{1u_n} & \cdots & x_{nu_n} \end{pmatrix} (u_0, c_0) < n$$

and

$$\begin{aligned} & \text{rank} \begin{pmatrix} x_{1u_1} & \cdots & x_{nu_1} & y_{u_1} & p_{1u_1} & \cdots & p_{nu_1} \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ x_{1u_n} & \cdots & x_{nu_n} & y_{u_n} & p_{1u_n} & \cdots & p_{nu_n} \end{pmatrix} (u_0, c_0) \\ &= \text{rank} \begin{pmatrix} x_{1u_1} & \cdots & x_{nu_1} & p_{1u_1} & \cdots & p_{nu_1} \\ \vdots & & \vdots & \vdots & & \vdots \\ x_{1u_n} & \cdots & x_{nu_n} & p_{1u_n} & \cdots & p_{nu_n} \end{pmatrix} (u_0, c_0) = n. \end{aligned} \quad (1)$$

Set $f(u, c) = \pi \circ \Gamma(u, c)$ and $\nu(u, c) = (-p(u, c), 1)/\sqrt{1 + |p(u, c)|^2}$. By a direct calculation, we have $f_{u_i}(u, c) \cdot \nu(u, c) = 0$ for all $(u, c) \in (\mathbb{R}^n \times \mathbb{R}^n, (u_0, c_0))$ and $i = 1, \dots, n$. It follows that (f, ν) is an n -parameter family of Legendre mappings (immersions). Moreover, we have

$$\begin{aligned} f_{c_i}(u, c) &= (x_{1c_i}(u, c), \dots, x_{nc_i}(u, c), y_{c_i}(u, c)) \\ &= \left(x_{1c_i}(u, c), \dots, x_{nc_i}(u, c), \right. \\ &\quad \left. \sum_{j=1}^n x_{jc_i}(u, c)g_{x_j}(x(u, c), p(u, c)) + \sum_{j=1}^n p_{jc_i}(u, c)g_{p_j}(x(u, c), p(u, c)) \right). \end{aligned}$$

It follows that $f_{c_i}(u, c) \cdot \nu(u, c) =$

$$\frac{1}{\sqrt{1 + |p(u, c)|^2}} \left(\sum_{j=1}^n (-p_j(u, c) + g_{x_j}(x(u, c), p(u, c))) x_{jc_i}(u, c) + \sum_{j=1}^n p_{jc_i}(u, c) g_{p_j}(x(u, c), p(u, c)) \right).$$

We now consider the following case. Suppose that

$$\text{rank} \begin{pmatrix} p_{1u_1} & \cdots & p_{nu_1} \\ \vdots & & \vdots \\ p_{1u_n} & \cdots & p_{nu_n} \end{pmatrix} (u_0, c_0) = n.$$

By using an n -parameter family of parameter change, we may assume that $p_i(u, c) = u_i$ for $i = 1, \dots, n$ by Proposition 3.11. If $e : (\mathbb{R}^n, \tilde{q}_0) \rightarrow (\mathbb{R}^n \times \mathbb{R}^n, (u_0, c_0))$ is a pre-envelope of (f, ν) , then

$$\begin{pmatrix} x_{1c_1}(e(q)) & \cdots & x_{1c_n}(e(q)) \\ \vdots & & \vdots \\ x_{nc_n}(e(q)) & \cdots & x_{nc_n}(e(q)) \end{pmatrix} \begin{pmatrix} (-p_1 + g_{x_1})(e(q)) \\ \vdots \\ (-p_n + g_{x_n})(e(q)) \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}.$$

Here we denote a local coordinate $(\mathbb{R}^n, \tilde{q}_0)$ by q instead of p in §3. Since Γ is an immersion germ and by Theorem 3.6, we have $(-p_i + g_{x_i})(e(q)) = 0$ for $i = 1, \dots, n$. It follows that $g_{p_i}(e(q)) = 0$ for $i = 1, \dots, n$ and hence $E(q) \in \pi(\Sigma_\pi(F)) = \pi(\Sigma_c(F))$. Conversely, if $E(q) \in \pi(\Sigma_c(F))$, then $f_{c_i}(e(q)) \cdot \nu(e(q)) = 0$ for all $i = 1, \dots, n$. By Theorem 3.6, e is a pre-envelope of (f, ν) .

Moreover, suppose that

$$\text{rank} \begin{pmatrix} x_{1u_1} & \cdots & x_{nu_k} & p_{k+1u_1} & \cdots & p_{nu_1} \\ \vdots & & \vdots & \vdots & & \vdots \\ x_{1u_n} & \cdots & x_{nu_k} & p_{k+1u_n} & \cdots & p_{nu_n} \end{pmatrix} (u_0, c_0) = n.$$

By using an n -parameter family of parameter change, we may assume that $x_i(u, c) = u_i$ for $i = 1, \dots, k$ and $p_j(u, c) = u_j$ for $j = k+1, \dots, n$ by Proposition 3.11. Then we also have $(-p_i + g_{x_i})(e(q)) = 0$ for $i = 1, \dots, k$ and $g_{p_j}(e(q)) = 0$ for $j = k+1, \dots, n$. It follows that $g_{p_i}(e(q)) = 0$ for $i = 1, \dots, k$ and hence $E(q) \in \pi(\Sigma_\pi(F)) = \pi(\Sigma_c(F))$. Conversely, if $E(q) \in \pi(\Sigma_c(F))$, then $f_{c_i}(e(q)) \cdot \nu(e(q)) = 0$ for all $i = 1, \dots, n$. By Theorem 3.6, e is a pre-envelope of (f, ν) .

The other cases, we can also prove by similarly. This completes the proof of Theorem. $\square$

By Theorems 4.1 and 4.2, if $\Sigma_c(F) = \Sigma_\pi(F)$ is an n -dimensional submanifold around z_0 , then $\Sigma_c(F)$ is a singular solution of $F = 0$ at z_0 and the projection $\pi(\Sigma_c(F))$ is an envelope when the variability condition holds.

We give concrete examples for completely integrable partial differential equations with a singular solution.

Example 4.3 Let $F : J^1(\mathbb{R}^2, \mathbb{R}) \rightarrow \mathbb{R}$ be given by $F(x_1, x_2, y, p_1, p_2) = -y + p_1^{n_1} + p_2^{n_2} = 0$, where $n_1, n_2 \geq 2$. That is, we consider the partial differential equation

$$y = \left(\frac{\partial y}{\partial x_1} \right)^{n_1} + \left(\frac{\partial y}{\partial x_2} \right)^{n_2}.$$

Then $\Sigma_c(F) = \Sigma_\pi(F) = \{(x_1, x_2, 0, 0, 0)\}$ is a 2-dimensional submanifold. The complete solution $\Gamma : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow F^{-1}(0)$ is given by

$$\Gamma(u_1, u_2, c_1, c_2) = \left(\frac{n_1}{n_1 - 1} u_1^{n_1 - 1} + c_1, \frac{n_2}{n_2 - 1} u_2^{n_2 - 1} + c_2, u_1^{n_1} + u_2^{n_2}, u_1, u_2 \right).$$

Then $(f, \nu) : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^3 \times S^2$ is a 2-parameter family of Legendre mappings, where

$$\begin{aligned} f(u_1, u_2, c_1, c_2) &= \pi \circ \Gamma(u_1, u_2, c_1, c_2) = \left(\frac{n_1}{n_1 - 1} u_1^{n_1 - 1} + c_1, \frac{n_2}{n_2 - 1} u_2^{n_2 - 1} + c_2, u_1^{n_1} + u_2^{n_2} \right), \\ \nu(u_1, u_2, c_1, c_2) &= \frac{1}{\sqrt{1 + u_1^2 + u_2^2}} (-u_1, -u_2, 1). \end{aligned}$$

Since

$$f_{c_i}(u_1, u_2, c_1, c_2) \cdot \nu(u_1, u_2, c_1, c_2) = -u_i / \sqrt{1 + u_1^2 + u_2^2}, \quad i = 1, 2,$$

$e : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \times \mathbb{R}^2, e(q_1, q_2) = (0, 0, q_1, q_2)$ is a pre-envelope and hence $E(q) = f \circ e(q) = (q_1, q_2, 0) \in \pi(\Sigma_c(F))$ is an envelope of (f, ν) .

Example 4.4 Let $F : J^1(\mathbb{R}^2, \mathbb{R}) \rightarrow \mathbb{R}$ be given by $F(x_1, x_2, y, p_1, p_2) = -y + p_1^{n_1} + x_2 p_2 + x_2^{n_2} = 0$, where $n_1, n_2 \geq 2$. That is, we consider the partial differential equation

$$y = \left(\frac{\partial y}{\partial x_1} \right)^{n_1} + x_2 \frac{\partial y}{\partial x_2} + x_2^{n_2}.$$

Then $\Sigma_c(F) = \Sigma_\pi(F) = \{(x_1, 0, 0, 0, p_2)\}$ is a 2-dimensional submanifold. The complete solution $\Gamma : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow F^{-1}(0)$ is given by

$$\Gamma(u_1, u_2, c_1, c_2) = \left(\frac{n_1}{n_1 - 1} u_1^{n_1 - 1} + c_1, u_2, u_1^{n_1} + \frac{2n_2 - 1}{n_2 - 1} u_2^{n_2} + c_2 u_2, u_1, \frac{n_2}{n_2 - 1} u_2^{n_2 - 1} + c_2 \right).$$

Then $(f, \nu) : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^3 \times S^2$ is a 2-parameter family of Legendre mappings, where

$$\begin{aligned} f(u_1, u_2, c_1, c_2) &= \pi \circ \Gamma(u_1, u_2, c_1, c_2) = \left(\frac{n_1}{n_1 - 1} u_1^{n_1 - 1} + c_1, u_2, u_1^{n_1} + \frac{2n_2 - 1}{n_2 - 1} u_2^{n_2} + c_2 u_2 \right), \\ \nu(u_1, u_2, c_1, c_2) &= \frac{1}{\sqrt{1 + u_1^2 + \left(\frac{n_2}{n_2 - 1} u_2^{n_2 - 1} + c_2 \right)^2}} \left(-u_1, -\frac{n_2}{n_2 - 1} u_2^{n_2 - 1} - c_2, 1 \right). \end{aligned}$$

Since

$$\begin{aligned} f_{c_1}(u_1, u_2, c_1, c_2) \cdot \nu(u_1, u_2, c_1, c_2) &= -\frac{u_1}{\sqrt{1 + u_1^2 + \left(\frac{n_2}{n_2 - 1} u_2^{n_2 - 1} + c_2 \right)^2}}, \\ f_{c_2}(u_1, u_2, c_1, c_2) \cdot \nu(u_1, u_2, c_1, c_2) &= \frac{u_2}{\sqrt{1 + u_1^2 + \left(\frac{n_2}{n_2 - 1} u_2^{n_2 - 1} + c_2 \right)^2}}, \end{aligned}$$

$e : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \times \mathbb{R}^2, e(q_1, q_2) = (0, 0, q_1, q_2)$ is a pre-envelope and hence $E(q) = f \circ e(q) = (q_1, 0, 0) \in \pi(\Sigma_c(F))$ is an envelope of (f, ν) .

Example 4.5 Let $F : J^1(\mathbb{R}^2, \mathbb{R}) \rightarrow \mathbb{R}$ be given by $F(x_1, x_2, y, p_1, p_2) = -y + p_1^{n_1} + x_2 p_2 + g(p_2) = 0$, where $n_1 \geq 2$ and g is a smooth function. That is, we consider the partial differential equation

$$y = \left(\frac{\partial y}{\partial x_1} \right)^{n_1} + x_2 \frac{\partial y}{\partial x_2} + g \left(\frac{\partial y}{\partial x_2} \right).$$

Then $\Sigma_c(F) = \Sigma_\pi(F) = \{(x_1, -g'(p_2), -g'(p_2)p_2 + g(p_2), 0, p_2)\}$ is a 2-dimensional submanifold. The complete solution $\Gamma : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow F^{-1}(0)$ is given by

$$\Gamma(u_1, u_2, c_1, c_2) = \left(\frac{n_1}{n_1 - 1} u_1^{n_1-1} + c_1, u_2, u_1^{n_1} + c_2 u_2 + g(c_2), u_1, c_2 \right).$$

Then $(f, \nu) : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}^3 \times S^2$ is a 2-parameter family of Legendre mappings, where

$$\begin{aligned} f(u_1, u_2, c_1, c_2) &= \pi \circ \Gamma(u_1, u_2, c_1, c_2) = \left(\frac{n_1}{n_1 - 1} u_1^{n_1-1} + c_1, u_2, u_1^{n_1} + c_2 u_2 + g(c_2) \right), \\ \nu(u_1, u_2, c_1, c_2) &= \frac{1}{\sqrt{1 + u_1^2 + c_2^2}} (-u_1, -c_2, 1). \end{aligned}$$

Since

$$\begin{aligned} f_{c_1}(u_1, u_2, c_1, c_2) \cdot \nu(u_1, u_2, c_1, c_2) &= -u_1 / \sqrt{1 + u_1^2 + c_2^2}, \\ f_{c_2}(u_1, u_2, c_1, c_2) \cdot \nu(u_1, u_2, c_1, c_2) &= (u_2 + g'(c_2)) / \sqrt{1 + u_1^2 + c_2^2}, \end{aligned}$$

$e : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \times \mathbb{R}^2$, $e(q_1, q_2) = (0, -g'(q_2), q_1, q_2)$ is a pre-envelope and hence $E(q) = f \circ e(q) = (q_1, -g'(q_2), -g'(q_2)q_2 + g(q_2)) \in \pi(\Sigma_c(F))$ is an envelope of (f, ν) .

References

- [1] V. I. Arnol'd, *Singularities of Caustics and Wave Fronts*. Mathematics and Its Applications **62** Kluwer Academic Publishers (1990).
- [2] V. I. Arnol'd, S. M. Gusein-Zade, A. N. Varchenko, *Singularities of Differentiable Maps vol. I*. Birkhäuser (1986).
- [3] J. W. Bruce, P. J. Giblin, What is an envelope? *Math. Gaz.* **65** (1981), 186–192.
- [4] J. W. Bruce, P. J. Giblin, *Curves and Singularities. A geometrical introduction to singularity theory. Second edition*. Cambridge University Press, Cambridge, 1992.
- [5] J. W. Bruce, P. J. Giblin, C. G. Gibson, Caustics through the looking glass. *Math. Intelligencer* **6** (1984), 47–58.
- [6] L. Chen, D. Pei, M. Takahashi, Envelopes of one-parameter families of frontals in hyperbolic and de-Sitter 2-space. *Preprint*. (2018).
- [7] S. Ei, K. Fujii, T. Kunihiro, Renormalization-group method for reduction of evolution equations; invariant manifolds and envelopes. *Ann. Physics.* **280** (2000), 236–298.
- [8] T. Fukunaga, M. Takahashi, Evolutes of fronts in the Euclidean plane. *J. Singul.* **10** (2014), 92–107.

- [9] T. Fukunaga, M. Takahashi, Evolutes and involutes of frontals in the Euclidean plane. *Demonstr. Math.* **48** (2015), 147–166.
- [10] T. Fukunaga, M. Takahashi, Framed surfaces in the Euclidean space. *Bull. Braz. Math. Soc. (N.S.)* (2018). DOI:10.1007/s00574-018-0090-z.
- [11] C. G. Gibson, *Elementary Geometry of Differentiable Curves. An undergraduate introduction.* Cambridge University Press, Cambridge, 2001.
- [12] A. Gray, E. Abbena, S. Salamon, *Modern differential geometry of curves and surfaces with Mathematica. Third edition.* Studies in Advanced Mathematics. Chapman and Hall/CRC, Boca Raton, FL, 2006
- [13] G. Ishikawa, *Singularities of Curves and Surfaces in Various Geometric Problems.* CAS Lecture Notes 10, Exact Sciences. 2015.
- [14] G. Ishikawa, Singularities of frontals. *Advanced Studies in Pure Mathematics.* **78** (2018), 55–106.
- [15] S. Izumiya, Singular solutions of first-order differential equations, *Bull. London Math. Soc.* **26** (1994), 69–74.
- [16] S. Izumiya, On Clairaut-type equations, *Publ. Math. Debrecen* **45** (1995), 159–166.
- [17] S. Izumiya, M. C. Romero-Fuster, M. A. S. Ruas, F. Tari, *Differential Geometry from a Singularity Theory Viewpoint.* World Scientific Pub. Co Inc. 2015.
- [18] T. Kunihiro, A geometrical formulation of the renormalization group method for global analysis. *Progr. Theoret. Phys.* **94** (1995), 503–514.
- [19] Y. Li, D. Pei, M. Takahashi, H. Yu, Envelopes of Legendre curves in the unit spherical bundle over the unit sphere. *Quarterly J. Math.* (2018) DOI: 10.1093/qmath/hax056.
- [20] L. Martins, K. Saji, Geometric invariants of cuspidal edges. *Canad. J. Math.* **68** (2016), 445–462.
- [21] M. Takahashi, On completely integrable first order ordinary differential equations, *Proceedings of the Australian-Japanese Workshop on Real and Complex singularities.* (2007), 388–418.
- [22] M. Takahashi, Envelopes of Legendre curves in the unit tangent bundle over the Euclidean plane. *Results Math.* **71** (2017), 1473–1489.
- [23] R. Thom, Sur la thorie des enveloppes. *J. Math. Pures Appl. (9)* **41** (1962), 177–192.

Masatomo Takahashi,
Muroran Institute of Technology, Muroran 050-8585, Japan,
E-mail address: masatomo@mmm.muroran-it.ac.jp