

Formulas of Frenet for a Vector Field in a Finsler Space

著者	NAGATA Yukiyoishi
journal or publication title	Memoirs of the Muroran Institute of Technology
volume	3
number	3
page range	543-546
year	1960-06-15
URL	http://hdl.handle.net/10258/3141

Formulas of Frenet for a Vector Field in a Finsler Space

Yukiyoshi Nagata

Abstract

T. K. Pan¹⁾ demonstrated the generalized formulas of Frenet for a vector field in a subspace of a Riemannian space. This paper extends his investigation to a hypersurface and a subspace of a Finsler space.

1. Formulas of Frenet for a Vector Field in a Hypersurface.

Let F_{n-1} be a hypersurface given by the set of equations $x^\lambda = x^\lambda(u^1, u^2, \dots, u^{n-1})$ ($\lambda=1, \dots, n$) in a Finsler space F_n the fundamental quadratic form of which is $ds^2 = g_{\lambda\mu}(x, x') dx^\lambda dx^\mu$. F_{n-1} to which the element of support is tangential has the fundamental quadratic form $ds^2 = {}_v g_{ab} du^a du^b$. Let v^λ be an arbitrary but fixed unit vector field defined at every point of F_{n-1} such that $v^\lambda = v^a B_a^\lambda$, ${}_v g_{ab} v^a v^b = 1$. Let $C: u^a = u^a(s)$ ($a=1, \dots, n-1$) be a curve on F_{n-1} and let N^λ be a unit vector normal to F_{n-1} . We define n vectors along C by the following equations:

$$\begin{aligned} \eta_{(1)}^\lambda &= v^\lambda, & \eta_{(2)}^\lambda &= {}_v k N^\lambda, \dots, \\ \eta_{(r+1)}^\lambda &= D\eta_{(r)}^\lambda / ds & (r &= 2, \dots, n-1), \end{aligned} \quad (1.1)$$

where ${}_v k = g_{\lambda\mu} N^\lambda Dv^\mu / ds$ and Dv^μ (${}^{2-3}$) denotes the absolute differential along C of the vector field v^μ at P of C . When $\eta_{(r)}^\lambda$ ($\beta=1, \dots, n$) are linearly independent, the following n vectors $\sigma_{(p)}^\lambda$ ($p=1, \dots, n$) which are expressed linearly with the components $\eta_{(r)}^\lambda$ for $r=1, \dots, p$ form a set of mutually orthogonal vectors:

$$\sigma_{(p)}^\lambda = \left(\frac{f_p}{f_{p-1}} \right)^{\frac{1}{2}} \eta_{(r)}^\lambda F_p^r \quad (r, \epsilon = 1, \dots, p) \quad (1.2)$$

where

$$\begin{aligned} f_0 &= 1, & f_1 &= 1, & f_p &= |f_r^\epsilon|, \\ f_r^\epsilon &= f_r^\epsilon = g_{\lambda\mu} \eta_{(r)}^\lambda \eta_{(\epsilon)}^\mu, & f_r^\epsilon F_p^r &= \delta_p^\epsilon. \end{aligned}$$

Putting

$$\frac{D\sigma_{(q)v}}{ds} \sigma_{(p)}^\nu = \alpha_{qp} \quad (p, q = 1, \dots, n), \quad (1.3)$$

from $\sigma_{(q)v} \sigma_{(p)}^\nu = \delta_q^p$ we have

$$-\alpha_{qp} = \alpha_{pq}, \quad (1.4)$$

$$\frac{D\sigma_{(p)}^\mu}{ds} = \sum_q \alpha_{pq} \sigma_{(q)}^\mu \tag{1.5}$$

From (1.1) and (1.2) it follows that $D\sigma_{(p)}^\mu/ds$ is at most a linear expression in $\eta_{(1)}^\mu, \dots, \eta_{(p+1)}^\mu$ and therefore in $\sigma_{(1)}^\mu, \dots, \sigma_{(p+1)}^\mu$. Consequently, $\alpha_{kh} = 0$ ($k+1 < h$). Combining this result with (1.4), we have

$$\begin{aligned} \alpha_{pp+1} &= -\alpha_{p+1p} = {}_vK_p \\ \alpha_{pq} &= 0 \quad (q \neq p \pm 1), \end{aligned} \tag{1.6}$$

where ${}_vK_p$ is defined by the first of these equations. Hence equations (1.5) are reduced to

$$\frac{D\sigma_{(p)}^\mu}{ds} = -{}_vK_{p-1} \sigma_{(p-1)}^\mu + {}_vK_p \sigma_{(p+1)}^\mu, \quad (p = 2, \dots, n-1), \tag{1.7}$$

where ${}_vK_p$ for $p=1, \dots, n-1$ are called, respectively, the associate curvatures of order $1, \dots, n-1$ of the vector field v for the curve C . (1.7) may be considered as a generalization of the Frenet formulas for a curve and hold except the case $p=1$. And (1.7) apply to the case $p=n$ with the understanding that ${}_vK_n=0$. We call these the *formulas of Frenet of the second kind for v along C in F_n* .

In the following, we shall derive the formulas of Frenet of the first kind for v along C in F_n . We put

$$\begin{aligned} \xi_{(1)}^\lambda &= v^\lambda, \quad \xi_{(2)}^\lambda = D\xi_{(1)}^\lambda/ds = {}_vK w^\lambda, \dots, \\ \xi_{(r+1)}^\lambda &= D\xi_{(r)}^\lambda/ds, \end{aligned} \tag{1.8}$$

where ${}_vK$ is the absolute curvature of v at P with respect to C and the sense of w is chosen in such a way as to make ${}_vK > 0$. If these vectors $\xi_{(\alpha)}^\lambda$ ($\alpha=1, \dots, n$) are assumed to be linearly independent, the following linear combinations of them for $p=1, \dots, n$ form a set of n mutually orthogonal vectors:

$$\mu_{(p)}^\lambda = \left(\frac{y_p}{y_{p-1}} \right)^{\frac{1}{2}} \xi_{(p)}^\lambda Y_p^\gamma \quad (\gamma, \epsilon = 1, \dots, p) \tag{1.9}$$

where

$$y_0 = 1, \quad y_p = |y_\epsilon^\epsilon|, \quad y_\epsilon^\epsilon = y_\epsilon^\epsilon = g_{\lambda\mu} \xi_{(p)}^\lambda \xi_{(\epsilon)}^\mu, \quad y_\epsilon^\epsilon Y_p^\epsilon = \delta_p^\epsilon.$$

And we have $\mu_{(1)}^\lambda = v^\lambda, \mu_{(2)}^\lambda = w^\lambda$.

Putting $(D\mu_{(h)}^\nu/ds) \mu_{(k)}^\nu = \beta_{hk}$ ($h, k=1, \dots, n$), from $\mu_{(h)}^\nu \mu_{(k)}^\nu = \delta_h^k$ we have

$$\beta_{kh} = -\beta_{hk} \tag{1.10}$$

$$\frac{D\mu_{(k)}^\nu}{ds} = \sum_h \beta_{kh} \mu_{(h)}^\nu. \tag{1.11}$$

(1.11) are reduced to

$$\frac{D\mu_{(k)}^\nu}{ds} = - {}_vL_{k-1}\mu_{(k-1)}^\nu + {}_vL_k\mu_{(k+1)}^\nu \quad (k = 2, \dots, n-1) \quad (1.12)$$

where ${}_vL_k = \beta_{kk+1} = -\beta_{k+1k}$.

(1.12) apply to the case $k=1$ with the understanding that ${}_vL_0=0$ and ${}_vL_1 = {}_vK$. Also, we have (1.12) for $k=n$ with the understanding that ${}_vL_n=0$. ${}_vL_k$ ($k=1, \dots, n-1$) are called, respectively, the associate curvatures of order $1, \dots, n-1$ of the vector field v for the curve C . We call (1.12) the *formulas of Frenet of the first kind for v along C in F_n* .

2. Extension.

We consider a subspace F_m ($m < n$) given by $x^\lambda = x^\lambda(u^1, \dots, u^m)$ ($\lambda=1, \dots, n$) in a Finsler space F_n . The element of support is tangential to F_m . Let N_p^λ ($p = m+1, \dots, n$) be $n-m$ mutually orthogonal unit vectors normal to F_m with respect to the metric of F_n . Let v^λ be an arbitrary but fixed unit vector field defined at every point of F_m such that $v^\lambda = v^a B_a^\lambda$, $g_{ab}v^a v^b = 1$ and $C: u^a = u^a(s)$ ($a=1, \dots, m$) be a curve on F_m .

We denote the absolute differential of v^λ with respect to C at P by Dv^λ and define the following vectors :

$$\begin{aligned} \eta_{(1)}^\lambda &= v^\lambda, & \eta_{(2)}^\lambda &= {}_v k_1 N_a^\lambda, \dots, \\ \eta_{(\gamma+1)}^\lambda &= \frac{D\eta_{(\gamma)}^\lambda}{ds} \quad (\gamma = 2, \dots, n-1), \end{aligned} \quad (2.1)$$

where ${}_v k_1 = g_{\lambda\mu} N_a^\lambda Dv^\mu / ds$.

If $\eta_{(p)}^\lambda$ ($\beta=1, \dots, n$) are linearly independent, the following linear combinations of them for $p=1, \dots, n$ form a set of n mutually orthogonal vectors :

$$\sigma_{(p)}^\lambda = \left(\frac{f_p}{f_{p-1}} \right)^{\frac{1}{2}} \eta_{(\gamma)}^\lambda F_p^\gamma \quad (\gamma, \epsilon = 1, \dots, p) \quad (2.2)$$

where

$$\begin{aligned} f_0 &= 1, & f_1 &= 1, & f_p &= |f_p^\epsilon|, \\ f_\gamma^\epsilon &= f_\epsilon^\gamma = g_{\lambda\mu} \eta_{(\gamma)}^\lambda \eta_{(\epsilon)}^\mu, & f_\gamma^\epsilon F_p^\gamma &= \delta_p^\epsilon. \end{aligned}$$

Therefore putting $(D\sigma_{(q)}^\nu / ds) \sigma_{(p)}^\nu = \alpha_{qp}$, we have

$$\frac{D\sigma_{(p)}^\mu}{ds} = - {}_vK_{p-1}\sigma_{(p-1)}^\mu + {}_vK_p\sigma_{(p+1)}^\mu \quad (p = 2, \dots, n-1), \quad (2.3)$$

where ${}_vK_p = \alpha_{pp+1} = -\alpha_{p+1p}$.

We call (2.3) the *formulas of Frenet of the second kind for v along C in F_n* . ${}_vK_p$ ($p=1, \dots, n-1$) are called, respectively, the associate curvatures of order $1, \dots, n-1$ of the vector field v for the curve C . (2.3) hold except the case $p=1$ and apply to the case $p=n$ with the understanding that ${}_vK_n=0$.

While, the formulas of Frenet of the first kind for v along C in F_n may be derived in the same way as is mentioned in 1.

(Received Jan. 26, 1960)

References

- 1) Pan, T. K.: Proc. Amer. Math. Soc. **8**, 294 (1957)
- 2) Davies, E. T.: Proc. London Math. Soc. **49**, 19 (1945)
- 3) Cartan, E.: Les espaces de Finsler P. 40 (Paris Hermann 1934)