

## Body composition and three indices of physical fitness in freshmen: exercise is of no effect?

著者	TANIGUCHI Kouji, SEINO Ichiji, KONARI Hidetoshi
journal or publication title	Memoirs of the Muroran Institute of Technology. Cultural science
volume	35
page range	141-148
year	1985-11-30
URL	<a href="http://hdl.handle.net/10258/1120">http://hdl.handle.net/10258/1120</a>

## Body composition and three indices of physical fitness in freshmen: exercise is of no effect?

著者	TANIGUCHI Kouji, SEINO Ichiji, KONARI Hidetoshi
journal or publication title	Memoirs of the Muroran Institute of Technology. Cultural science
volume	35
page range	141-148
year	1985-11-30
URL	<a href="http://hdl.handle.net/10258/1120">http://hdl.handle.net/10258/1120</a>

## Body composition and three indices of physical fitness in freshmen: exercise is of no effect ?

Kôji Taniguchi, Ichiji Seino, and Hidetoshi Konari.

### Abstract

Grip strength, back strength, standing trunk flexion and skinfold thickness were measured on 314 freshmen (mean 18.6 years). Body composition was estimated by the method of measuring skinfold thickness. The subjects were asked whether they had belonged to athletic clubs in junior and/or senior high schools or not, and they were divided into four groups with regard to their answers.

Of the groups, group A is composed of 94 students who belonged to athletic clubs both in junior and senior high schools and group B is composed of 117 students who did not. The values of groups A and B were compared.

The results were as follows:

- 1) The mean value of fat storage rate (% fat) of all subjects was 21% and 44 per cent of them were judged to be obese.
- 2) The % fat of group A was smaller than that of group B.
- 3) Group A indicated significantly bigger Lean Body Mass (LBM) and LBM/Height-index than group B ( $P < 0.01$ ).
- 4) The grip strength and back strength of group A were significantly better than those of group B ( $P < 0.01$ ).
- 5) There was no statistical difference in standing trunk flexion between groups A and B.

From these results, it seems that exercise in adolescence keeps % fat small, promotes the development of muscle and enhances muscular strength.

The development of materialistic civilization have caused the shortage of physical activities. The shortage of physical activities is apt to induce obesity<sup>1), 2)</sup> and various diseases.<sup>3), 4)</sup> How should we judge obesity ? The method of judging obesity from the index of height and weight have been widely used: Rohrer index, Ponderal index, Kaup index, etc.<sup>5), 6)</sup> Suppose there are two persons of the same height and weight: one leads an active life, the other does

not. Can we conclude that there is no difference in body between the two persons ?

Generally speaking, endurance runners have smaller adipose tissue and weight lifters have bigger muscular tissue than inactive persons.<sup>7),8)</sup> Judging obesity by the indices of anthropometry could be misleading. That is the reason why we apply the conception of "body composition". Body composition can be divided into Lean Body Mass and adipose tissue.<sup>9),10),11)</sup> Body composition can not be directly measured *in vivo*, and so it is indirectly estimated various methods.<sup>10),12)~17)</sup> Of the methods, the method of measuring skinfold thickness is widely applied because of its simplicity.<sup>6),18),19),20)</sup>

There are many papers on the relation of physical activities to body composition.<sup>7),8),21)~24)</sup> What about body composition in Japanese freshmen ? Masaki<sup>25)</sup> reported that the values of back strength and standing trunk flexion in Japanese pupils were decreasing. What about those in Japanese freshmen ?

The purpose of this paper is to estimate body composition in Japanese freshmen, and to examine the effect of exercise on body composition and three indices of physical fitness.

## Methods

### Subjects

Three hundred and fourteen freshmen in Muroran Institute of Technology participated in this study as subjects. They ranged in age from 18 to 21 years.

### Body composition (% fat, LBM)

Skinfold thickness was measured with "Eikenshiki skinfold caliper".<sup>26)</sup> % fat was estimated by the modified method of Nagamine<sup>18)</sup> and that of Brožek *et al.*<sup>27)</sup> LBM was calculated by using % fat derived from the methods described above:  $LBM = (1 - \% \text{ fat}/100) \times \text{Weight}$ . LBM represents muscular mass.<sup>11)</sup> LBM/Height-index represents the development of muscle.<sup>11)</sup> % weight was calculated

Body composition and three indices of physical fitness in freshmen : exercise is of no effect ?

on the basis of the standard weight in the hand book published from the Ministry of Welfare.<sup>28)</sup>

## Results

Physical characteristics and measurements of subjects are presented in table1. We compared our results with statistical data.<sup>29),30)</sup> Our subjects were similar to those in the data in height, but significantly heavier in weight. Back strength and standing trunk flexion were inferior to those in the data, but grip strength was better. Skinfold thickness in tricep showed no significant difference, but that of subscapular showed significant difference.

**Table 1** Physical characteristics and measurements of subjects. N=314

	Height (cm)	Weight (kg)	Rohrer index	% weight	S.T. tricep	S.T. subs.	S.T. sum
m	170.6	63.6	128.3	107.4	13.6	17.1	30.7
SD.	5.45	8.98**	18.12	14.35	7.22	9.45**	16.00**
	% fat	LBM	LBM/Height	Grip S.	Back-S.	Standing T.F.	
m	21.0	49.8	29.1	48.4	92.6	13.2	
SD.	7.87	5.02	2.46	7.16**	19.62**	5.99**	

S.T.: skinfold thickness. sum: tricep + subscapular.

\* and \*\* show the significance at 5% and 1% level, respectively.

Nagamine<sup>6)</sup> and Huenemann, R. L., *et al.*<sup>2)</sup> recommend 20% fat as the criterion of obesity. In this study, 44 percent of all subjects were considered to be obese. LBM and LBM/Height-index showed smaller values than those of other reports.<sup>31)~34)</sup>

Table 2 Comparison of the values between groups A and B.

		Height (cm)	Weight (kg)	Rohrer index	% weight	S.T. tricep	S.T. subs.	S.T. sum.
A: n=94	m	171.4	64.9	129.2	108.7	13.6	16.2	29.8
	S.D	5.22	7.96	17.03	13.14	7.08	8.61	14.99
B: n=117	m	169.5	62.3	127.9	106.4	14.5	18.4	32.9
	S.D	5.17**	9.17*	18.63	15.18	7.79	10.35	17.5
		% fat	LBM	LBM/Height	Grip S.	Back S	Standing T.F.	
A: n=94	m	20.4	51.3	29.9	49.2	95.9	13.7	
	S.D	7.37	4.80	2.30	6.87	18.36	6.51	
B: n=117	m	22.3	47.8	28.2	46.4	87.2	12.6	
	S.D	8.66	4.65***	2.35***	7.08***	20.35***	5.76	

Table 2 shows comparison of group A with group B. Height and weight showed significant difference ( $P < 0.01$ ,  $P < 0.05$ ). Skinfold thickness and % fat in group B showed large values. Muscular strengths were better in group A. LBM and LBM/Height-index were better in group A, too.

Table 3 shows the distribution of the degree of obesity from the viewpoints of % weight and % fat, Thirty per cent of subjects were judged to be obese with regard to % fat, but they could not be judged to be obese with regard to % weight. This tendency can be seen more in group B than in group A.

**Table 3** Distribution of the degree of obesity from the viewpoints of % weight and % fat

% fat		% weight	non-obesity				obesity
			~90	91~100	101~110	111~120	121~
non-obesity	~14		4	12	5	1	
		middle	7	15	4		
		under	19	40	15	1	
	15~19			11	14	7	
		middle	4	21	4	2	
		under	6	43	37	14	
obesity	20~24		2	11	5	1	
		middle	4	14	6	1	
		under		7	36	18	5
	25~29			1	5	6	
		middle		7	6	1	
		under		8	15	11	
	30~				3	6	
		middle			7	14	
		under			12	27	

above column; numbers of group A.  
middle column; numbers of group B.  
under column; numbers of all subjects.

### Consideration

The shortage of physical activities may replace a part of muscle with adipose.<sup>35)</sup> In this study, mean values of Rohrer index and % weight show no obesity, but those of % fat show obesity. Consequently we might say that many students wear excessive adipose coats.<sup>1)</sup> In other words, there may be many students lacking in sufficient muscular mass. This tendency is apt to be seen in group B.

Although Masaki<sup>25)</sup> reported that muscular strength was decreasing, we con-

sider this tendency as an excessive adiposity or lack of sufficient muscular mass. The moderns seem to have inactive life-styles. We expect that the majority of our subjects who have little experience of walking a long distance and exerting all their powers will have undeveloped muscles. This seems to be supported by the comparison between groups A and B: in group A, % fat was smaller, muscular strengths were better, LBM and LBM/Height-index were higher. We think that these are the effect of exercise in adolescence.

Although we have gained such conclusion as above, we concern about the results in group A. % fat of group A was higher than that of other reports. In addition LBM and LBM/Height-index were smaller than those in other reports.<sup>32), 38)</sup> The values of group A were similar to those of inactive American young men.<sup>39)</sup> We must consider that the inactive life of students preparing for entrance examination had a great influence on our subjects.

Finally, we intend to increase the subjects of our investigation. We will get more information on body composition as well as on exact physical activities and make training programs for individuals.

(昭和60年5月21日 受理)

## References

- 1) Stefanik, P. A., *et al.*, "Caloric intake in relation to energy output of obese and non-obese adolescent boys," *Am. J. Clin. Nutrition*, 7 : 55-62, 1959.
- 2) Huenemann, R. L., *et al.*, "Adolescent food practices associated with obesity," *Fed. Proc.*, 25 : 4-10, 1966.
- 3) Keys, A., *et al.*, "Coronary heart disease: overweight and obesity as risk factors," *Ann. Int. Med.*, 77 : 15-27, 1972.
- 4) クラウス・ラーブ (広田公一・石川亘訳), 運動不足病, ベースボール・マガジン社, 1977. Pp. 230. (Kraus, H. and Raab, W., *Hypokinetic disease: diseases produced by lack of exercise*, Thomas Co.: Springfield, 1961.)
- 5) Keys, A. *et al.*, "Indices of relative weight and obesity," *J. Chron. Dis.*, 25 : 329-43, 1972.
- 6) 長嶺晋吉「肥満の判定法」*医学のあゆみ*, 101 : 404-09, 1977.



Body composition and three indices of physical fitness in freshmen : exercise is of no effect ?

- 7) Boileau, R. A. *et al.*, "Body composition changes in adult men during selected weight training and jogging programs," *The Research Quarterly*, 44 : 158—68, 1963.
- 8) Pollock, M. L., *et al.*, "Body composition of elite class distance runners," *Ann. N. Y. Acad. Sci.*, 301 : 361—70, 1977.
- 9) Behnke, A. R., *et al.*, "The specific gravity of healthy men: body weight  $\div$  volume as an index of obesity," *J. A. M. A.*, 118 : 495—98, 1942.
- 10) Pace, N. and Rathbun, E. N., "Studies on body composition III . The body water and chemically combined nitrogen content in relation to fat content," *J. Biol. Chem.*, 158 : 685—91, 1945.
- 11) Behnke, A. R., *et al.*, "Lean body mass: Its clinical significance and estimation from excess fat and total body water determinations," *A. M. A. Arch. Intern. Med.*, 91 : 585—601, 1953.
- 12) Ward, A., *et al.*, "A comparison of body fat determined by underwater weighing and volume displacement," *Am. J. Physiol.*, 234 : E94—E96, 1978.
- 13) Kyriakopoulos, V., *et al.*, "The radiological assessment of body fat," *Clin. Radiol.*, 30 : 477—79, 1979.
- 14) Forbes, G. B. and Hursh, J. B., "Age and sex trends in lean body mass calculated from  $K^{40}$  measurements: with a note on the theoretical basis for the procedure," *Ann. N. Y. Acad. Sci.*, 110 : 255—63, 1963.
- 15) Lesser, G. T., *et al.*, "Determination of total body fat by absorption of an inert gas: measurements and results in normal human subjects," *J. Clin. Invest.*, 39 : 1791—1806, 1960.
- 16) Pollock, M. L., *et al.*, "Prediction of body density in young and middle-aged men," *J. Appl. Physiol.*, 40 : 300—04, 1976.
- 17) Wilmore, J. H., *et al.*, "Validity of skinfold and girth assessment for predicting alterations in body composition," *J. Appl. Physiol.*, 29 : 313—17, 1970.
- 18) Brožek, J., *et al.*, "Skinfold distributions in middle-aged American men: a contribution to norms of leanness—fatness," *Ann. N. Y. Acad. Sci.*, 110 : 492—502, 1963.
- 19) Durnin, J. V. G. A. and Rahaman, M. M., "The assessment of the amount of fat in the human body from measurements of skinfold thickness," *Br. J. Nutr.*, 21 : 681—89, 1967.
- 20) Lohman, T. G., "Skinfolds and body density and their relation to body fatness: a review," *Human Biol.*, 53 : 181—225, 1981.
- 21) 長嶺晋吉他「スポーツマンと非スポーツマンの体構成 (Body Composition) の比較に関する研究」*栄養学雑誌*, 24 : 3—8, 1965.
- 22) Lewis, S., *et al.*, "Prediction of body composition in habitually active middle-aged

- men," *J. Appl. Physiol.*, 39 : 221-25, 1975.
- 23) Moore, C. E., *et al.*, "The relationship of exercise and diet on high-density lipoprotein cholesterol levels in women," *Metab.*, 32 : 189-96, 1983.
- 24) Pařízková, *et al.*, "Body composition, aerobic capacity, and density of muscle capillaries in young and old men," *J. Appl. Physiol.*, 31 : 323-25, 1971.
- 25) 正木健雄, 子どもの体力, 大月書店, 1979. Pp 214.
- 26) 長嶺晋吉「皮厚の測り方と意義」 臨床医, 2 : 538-40, 1976.
- 27) Brožek, J., *et al.*, "Densitometric analysis of body composition: revision of some quantitative assumptions," *Ann. N. Y. Acad. Sci.*, 110 : 113-40, 1963.
- 28) 長嶺晋吉, 「肥満の判定法」, 鈴木慎次郎・野村茂 (編), 生活と肥満, 医歯薬出版, 1981. pp 50-51.
- 29) 厚生省栄養課編, 国民栄養の現状, 第一出版, 1983. Pp 173.
- 30) 東京都立大学身体適性学研究室編, 日本人の体力標準値, 第3版, 不昧堂出版, 1980. Pp458.
- 31) 池田修他「福岡市とその近郊の中学生及び大学生(年齢12. 13歳及び18. 19歳)の体組成と有酸素能力の関係」 人類学雑誌, 86 : 363-66, 1978.
- 32) 北川薫他「最大酸素摂取量の規定因子としての除脂肪体重の検討」 体力科学, 23 : 96-100, 1974.
- 33) 今野道勝他「福岡市近郊の成人男女の栄養・運動・身体組成について」 健康科学, 3 : 97-104, 1981.
- 34) 北川薫「成人男女の身体組成とその国際比較」 中京体育学研究, 18 : 51-56, 1977.
- 35) Sandler, H., "Use of bed rest studies to simulate the problems of human existence in a space flight environment," Ames. Res. Center, NASA, USA. (cited by Ikegami: 運動処方, 朝倉書店, 1982. p39)
- 36) Komiya, S., and Kikkawa, K., "Change in body composition of Japanese children and adolescents during the last 16 years," *J. Health Science*, 1 : 15-26, 1979.
- 37) Komiya, S. and Kikkawa, K., "Hand-grip strength and standing long jump of Japanese children of two different cohorts in relation to their body composition," *J. Health Science*, 2 : 49-56, 1980.
- 38) 増田卓二・吉水浩「日常生活における身体活動レベルと最大酸素摂取量( $\dot{V}O_2\text{max}$ )の関係(ネパール人を対象とした研究調査報告)」 久留米大学論叢, 28 : 61-65, 1979.
- 39) Womersley, J. *et al.*, "Influence of muscular development, obesity, and age on the fat-free mass of adults," *J. Appl. Physiol.*, 41 : 223-29, 1976.